

Preliminary phylogeny of *Tribolium* beetles (Coleoptera: Tenebrionidae) resolved by combined analysis of mitochondrial genes

NEVENKA MEŠTROVIĆ, BRANKICA MRAVINAC, MIROSLAV PLOHL, ĐURĐICA UGARKOVIĆ and BRANKA BRUVO-MAĐARIĆ*

Ruder Bošković Institute, Bijenička 54, HR-10002 Zagreb, Croatia

Key words. *Tribolium*, mitochondrial DNA, cytochrome oxidase I, 16S rDNA, phylogeny, sibling species

Abstract. The phylogenetic relationships of the three major species groups of *Tribolium* (Coleoptera: Tenebrionidae) were inferred using the simultaneous analysis of 642 bp of the most conserved part of mitochondrial DNA (mt DNA) cytochrome oxidase I (*COI*) and 448–452 bp of mt 16S rDNA. High sequence divergence was observed for both genes even among sibling species. The analysis of the combined segments of *COI* and 16S rDNA sequences produced a phylogenetic tree with moderate level of confidence. The tree topology showed monophyly of the genus *Tribolium* whose species were separated into three groups: “brevicornis” group (with *T. brevicornis* as the only representative), “castaneum” group (with *T. castaneum*, *T. freemani*, *T. madens* and *T. audax*) and “confusum” group (with *T. confusum*, *T. anaphe* and *T. destructor*). Sibling species pairs *T. castaneum* – *T. freemani* and *T. madens* – *T. audax* are clearly resolved. The preliminary results presented here give moderate support to the previously proposed phylogeny based on morphological data.

INTRODUCTION

The genus *Tribolium* Macleay, 1825 belongs to the family Tenebrionidae (order Coleoptera) and comprises 33 described species (Sokoloff, 1972), several of which are important stored product pests. Some of the species are easily cultured in the laboratory and the genus has been extensively used in genetic analyses (Sokoloff, 1972; Beeman et al., 1996; Bennett et al., 1999; Brown et al., 2000). Thanks to its convenient genetic set-up, *Tribolium castaneum* is predisposed to become one of the three most important insect model systems beside *Drosophila* and *Bombyx mori* (Schmidt & Tautz, 1999; Brown et al., 2003; Nagaraju & Goldsmith, 2002).

Several studies have attempted to establish evolutionary trends in these beetles. According to morphological characters, Hinton (1948) divided *Tribolium* species into five groups: “brevicornis” group (considered as the most primitive), “confusum”, “alcine”, “castaneum” and “myrmecophilum” group. According to Hinton, the first four groups separated in the Cretaceous period, while the “myrmecophilum” group separated from the “castaneum” group in the Quaternary. The “brevicornis”, “castaneum” and “confusum” species-groups are predominant in the number of species and their distribution, and can be further split into sections. The remaining two groups comprise only five species that are geographically restricted to Australia and Madagascar. Within *Tribolium*, even the sibling species *T. madens* – *T. audax* and *T. castaneum* – *T. freemani*, able to hybridize, are morphologically distinguishable by numerous characters pointing to a high species divergence in the group (Halstead, 1969; Wade & Johnson 1994).

Cladistic relationships among some *Tribolium* species inferred from cytological studies of chromosome numbers, revealed the “castaneum” group as more “primitive” than the “confusum” group (Smith, 1952). Evolutionary relationships of the most common *Tribolium* species were also studied by electrophoretic analysis of isozymes (Wool, 1982). According to this study, *T. castaneum* and *T. confusum* are more closely related to *T. brevicornis* than they are to each other, which is in

disagreement with cytological and morphological data. A comparison of the defensive secretions from *Tribolium* species indicated that “brevicornis” and “castaneum” are sister species groups, and that *T. brevicornis* and *T. audax* are members of the same species group (Howard, 1987).

To our knowledge, there has been no study of this genus so far that would be based on phylogenetically informative molecular data. At the DNA level, relationships between some *Tribolium* species were inferred using characteristics of highly repetitive DNAs, their sequence homology and abundance (Juan et al., 1993; Ugarković et al., 1996). The cladogram obtained from satellite DNA hybridization data, karyological data and chemical compounds of secretion (Juan et al., 1993) supported the “castaneum” species-group, although *T. audax* was placed outside of this group, while “confusum” and “brevicornis” species-groups appeared as sister clades, in contrast to the phylogeny proposed by Hinton (1948) and Howard (1987).

In this work we address the phylogenetic relationships among the eight most studied species of *Tribolium*: *T. castaneum*, *T. confusum*, *T. destructor*, *T. madens*, *T. audax*, *T. anaphe*, *T. freemani* and *T. brevicornis*, which are also the most widespread and economically important. These eight species represent, according to Hinton (1948), three major species groups: the “castaneum” group (*T. castaneum*, *T. audax*, *T. freemani* and *T. madens*), the “confusum” group (*T. confusum*, *T. anaphe*, *T. destructor*) and the “brevicornis” group (*T. brevicornis*). Despite our efforts, we were unable to get hold of any of the additional “brevicornis” group species, nor any of the species belonging to “alcine” and “myrmecophilum” groups, and therefore we regard this study as an initial and preliminary analysis of *Tribolium* phylogeny. In order to resolve the relationships of these eight species we used two genes that are regarded as the most conserved among mitochondrial markers routinely used for phylogenetic inference: cytochrome oxidase I (*COI*) and 16S rDNA. *COI* has been used for phylogenetic inference of many insects (Lunt et al., 1996) – the variable part of this gene has been used to deduce phylogenetic relationships among congeneric species

* Corresponding author; e-mail: bruvo@irb.hr

of the tenebrionid genera *Pimelia* and *Hegeter* (Juan et al., 1995, 1996; Pons et al., 2004). However, this part of the *COI* gene appeared to be too variable for phylogenetic reconstruction within the tenebrionid genus *Palorus* (Meštrović et al., 2000), a taxon closely related to the genus *Tribolium*. For this reason, we decided to use the most conserved part of the *COI* gene, proposed by Zhang & Hewitt (1997) to be suitable for low- and mid level phylogenetic analysis. The other phylogenetic marker used in this study, 16S rDNA, appears to be at least twice as conserved as *COI* gene in species of the order Coleoptera (Funk, 1999; Garin et al., 1999).

MATERIAL AND METHODS

Beetle material

Insect cultures (laboratory strains) used in this study were obtained from the Central Science Laboratory (Sand Hutton, York, UK). A natural population of the species *T. madens* was collected in a flourmill (Virovitica, northern Croatia). Total DNA was extracted from a single individual according to Steiner et al. (1995), or from pooled individuals following the standard protocols.

PCR amplification and sequencing

Parts of *COI* were amplified using primers C1-J-2090 (UEA5) (5'AGTTTTCAGCAGGAGCAATTACTAT3') and C1-N-2738 (UEA8) (5'AAAAATGTTGAGGGAAAAATGTTA3') (Zhang & Hewitt, 1997). These primers amplify about 690 bp of the *COI* gene, corresponding to sites 1747 and 2392 of the *Drosophila yakuba* mitochondrial genome. Primers LR-N-13388 (5'CGCCTGTTTATCAAAAACAT3') and a modified LR-J-12887 (5' CTCCGGTTTGAACCTCAGATCA3') (Simon et al., 1994) were used to amplify a segment of about 550 bp of the mitochondrial 16S ribosomal DNA. Sequences of 16S rDNA from *T. madens* were amplified with internal primers (LR-J1: 5'GTAAAATTTTAAAGGTGCAACAGACC3' and LR-N1: 5'ATTATTATTTAAGGTGCAACCTGCCC3'). The reaction conditions were the following: 1 × Taq buffer, 2.5 mM MgCl₂, 200 μM of each dNTP, 0.4 μM of each primer and 1U of Taq polymerase (Promega, Southampton, UK). Thermocycling consisted of an initial denaturation step of 3 min at 95°C followed by 35 cycles of denaturation for 30 s at 95°C, annealing for 1 min at 50°C, extension for 1.5 min at 72°C and finally, a single extra extension step of 10 min at 72°C (thermocycler Perkin Elmer 2400). PCR products were purified with GFX PCR purification Kit (Amersham-Pharmacia, Buckinghamshire, UK) and sequenced either directly or after cloning into the pGEM-T vector (Promega) for the PCR products obtained from "pooled" DNA specimens. Sequencing was carried out in both directions by a DNA sequencing Service (MWG-Biotech, Ebersberg, Germany). Sequences have been deposited in the EMBL databank under accession numbers AJ438061 – AJ438096 and DQ304692 – DQ304693 for *COI*, and AJ438140 – AJ438154 and AJ566071 for 16S.

Phylogenetic analyses

The alignment of the *COI* and 16S rDNA sequences was done with CLUSTAL X version 1.81 (Thompson et al., 1997). BioEdit 5.09 (Hall, 1999) was used for manual inspection and correction of 16S rRNA alignments according to secondary structure of *Drosophila yakuba* 16S rRNA (Gutell & Fox, 1988). Aligned sequences were reformatted appropriately and entered into MEGA 2.1 (Kumar et al., 2001) for calculation of sequence statistics. In order to downweight the gaps present in 16S ribosomal gene (24 gaped positions altogether), we applied the "simple method" for recoding gaps described in Simmons & Ochoterena (2000) and Simmons et al. (2001) using the program

GapCoder (Young & Healy, 2002). The matrices obtained this way were then used in both parsimony and Bayesian searches. The alignments used in phylogenetic analyses are available on request from the authors.

COI and 16S rDNA data sets were tested using the partition homogeneity test (Farris et al., 1994) implemented within PAUP to assess whether combining of *COI* and 16S sequences is appropriate for phylogenetic analysis. Phylogenetic analyses were performed on separate and combined datasets by parsimony (MP) and maximum likelihood (ML) searches using PAUP 4.0b10 (Swofford, 1998), and by Bayesian analysis with Markov Chain Monte Carlo (MCMC) sampling in MrBayes 3.1 (Huelsenbeck & Ronquist, 2001).

For MP analyses, we used the heuristic search with tree-bisection-reconnection (TBR) branch-swapping option. The extent of support for internal nodes was estimated using the bootstrap method (1000 pseudoreplicates) (Felsenstein, 1985). Bremer support and partitioned Bremer support values (Baker & DeSalle, 1997) were calculated based on strict consensus MP trees using the program TreeRot 2 (Sorenson, 1999).

The model of DNA substitution was determined for each gene with the program MODELTEST 3.04 (Posada & Crandall, 1998). The estimated parameters (GTR+G+I, being the model preferred by both Akaike information criterion (AIC) and hierarchical likelihood ratio test (hLRT) for both partitions) were then invoked for ML heuristic searches with simple sequence addition and TBR branch-swapping option.

Bayesian MCMC analyses with four parallel Markov chains were run under the GTR+G+I model (unlinked parameters for each partition, overall rate allowed to be different across partitions) for 1000000 generations, with sampling frequency of one in every hundred trees. The consensus tree (with posterior probabilities for each node as a measure for support) was constructed based on the trees sampled after the convergence of likelihood scores (burn-in).

RESULTS

Sequence variation of *COI* gene

Sequence analysis of a 642 bp fragment of the most conserved part of mitochondrial *COI* gene was performed in eight *Tribolium* species as well as in two outgroup species, *Pimelia elevata* and *Tenebrio molitor*. Intraspecific variation, assessed by screening four to eight specimens of each species, ranges from 0 to 1%, with the exception of *T. freemani* where intraspecific variability is 2.3% mainly due to the slippage in one of the four analysed clones. In addition, one of *T. brevicornis* sequences appeared shorter due to several internal deletions leading to frame-shift mutations. It can be suspected that such a sequence represents a nuclear paralogue of a part of *COI* gene, and for this reason it was not considered for further analyses. Other *COI* fragments did not present any peculiarities that would point to their nuclear provenience (review on mitochondrial pseudogenes in Bensasson et al., 2001). In the case of *T. madens*, the specimens from two different populations, laboratory and natural, were analyzed. Differences between these populations (0.06–0.50%) are within the range of intraspecies divergences of other examined species.

Among the 642 examined nucleotide positions of tested *Tribolium* sequences, 223 positions were polymorphic (34.7%) and 153 (23.8%) phylogenetically informative. Nucleotide composition, number of variable sites and Ti/Tv ratios are presented in Table 1. The sequences have an average frequency of A+T nucleotides of 63.0%, increasing to 74.3% at third codon positions. Similar frequency of A+T has been observed in two other tenebrionid genera – *Pimelia* (Juan et al., 1995) and *Hegeter*

(Juan et al., 1996). The observed A+T bias is lower than in other insects, what is suggested to be a general characteristic of beetle *COI/COII* genes (Liu & Beckenbach, 1992; Funk et al., 1995). The average transition/transversion ratio (calculated from the alignment of the entire dataset) in examined sequences is 1.3, being the highest at first codon positions, while dropping close to 1 at third codon positions. Of the 223 variable positions, 80.7% are at third codon positions, 15.7% at first, while the second codon positions are the most conserved, with only 3.6% of nucleotide changes (Table 1).

Comparison of inferred amino acid (aa) sequences of COI in *Tribolium* using mtDNA code for *D. melanogaster* (De Bruijn, 1983) showed that the 214 aa long polypeptide is highly conserved with only 20 changes (10%) of which 12 are conservative. Most of these changes are located in the variable part of the protein between positions 182–212, what is in accordance with the suggested aa variability of insect COI (Lunt et al., 1996). Maximum aa divergence between pairs of sequences within *Tribolium* amounts to 7%.

Uncorrected p-distances were calculated for each species pair. The minimum distance of 12% is calculated for *T. confusum*–*T. anaphe* species pair, while all other sequence pairs are within the range from 14% to the maximum divergence of 19% for *T. destructor*–*T. audax*. The genetic distances between *Tribolium* and the outgroups (*Tenebrio molitor* and *Pimelia monticola*) range between 16% and 20%. The fact that some ingroup divergences are higher than the ingroup-outgroup distance, together with low T_s/T_v ratio, and high number of T_v A↔C as well as variable sites in the 3rd codon position (80.7%), suggests that there is a high level of saturation due to multiple changes in the *COI* data set.

Sequence variation of 16S rDNA

Sequence analysis of a 448–452 bp fragment of the 16S rDNA has been performed in the same species used to obtain *COI* data. Intraspecific variation was in the range from 0 to 0.89%. The fragment length was conserved within the species, but in the alignment of *T. freemani* clones two gaps are created, although the clones have the same length. Since analyzed 16S rDNA sequences showed unexpectedly high interspecific divergence (see below), they could not be easily aligned with a significant confidence. The computer-generated alignment was therefore further adjusted considering published secondary structures for *Drosophila yakuba* 16S rRNA (Gutell & Fox, 1988). Some stems show noncanonical G-T pairs, which also help in maintaining stem stability (Gutell et al., 1994). In the highly variable region between positions 190–249, which corresponds to G3 helix (De Rijk et al., 1996), it was not possible to predict the stem and loop regions.

Sequence analysis reveals significantly higher overall A+T content in 16S than in *COI* (Table 1). The base composition is clearly biased towards AT, especially in the loops. The percent of variable sites is similar between stems and loops, consistent with previous observations that both regions contain sites allowed to evolve with heterogeneous rates regardless of their secondary structure (Simon et al., 1994). If G3 helix is excluded from the analysis, the percentage of variable sites is lower in 16S than in *COI*, but when G3 is considered, the divergences are highly similar. Since it is expected that insect 16S rDNA evolves at least twice slower than *COI* (Funk, 1999), this suggests saturation of 16S rRNA gene as well. The overall T_s/T_v ratio of 16S is twice as low as that of *COI*, as expected for 16S and *COI* sequences showing similar levels of divergence (Funk et al., 1995). The 16S T_s/T_v ratio was twice higher in stems (similar to that of *COI*) than in loops, in which T_s may be completely saturated.

TABLE 1. Summary of nucleotide composition, number of variable sites and T_s/T_v ratio for *COI* and 16S rDNA data of examined *Tribolium* beetles.

	base pairs	nucleotide %					variable sites	Ts/Tv ratio
		A	T	C	G	A+T		
<i>COI</i>								
all	642	31.6	31.4	20.5	16.4	63.0	223	1.3
1st pos	214	29.1	22.9	18.1	29.7	52.0	35	3.0
2nd pos	214	20.2	42.1	21.6	15.9	62.3	8	1.7
3rd pos	214	45.1	29.2	21.7	3.5	74.3	180	1.1
16S								
all	448–452	32.6	40.1	8.0	19.3	72.7	148	0.61
stem *	209	38.1	31.0	8.1	22.7	69.1	51	1.01
loop *	182–185	34.7	38.9	9.9	16.5	73.6	51	0.54

*exclusion of G3 helix (between position 190–249) T_s/T_v , transition/transversion ratio = (no. inferred T_s)/(no. inferred T_v).

Within the examined *Tribolium* species, genetic distances reveal the minimum value of 7% for the *T. madens* – *T. audax* pair and 9% for the *T. anaphe* – *T. confusum* pair, and the maximum value of 18% for the *T. freemani* – *T. destructor* pair. Genetic distances between *Tribolium* and the *Tenebrio* / *Pimelia* outgroups are in the range from 16 to 22%.

Phylogenetic analysis

COI

COI trees obtained by all three methods revealed the same topology (not shown). Each of the species is clearly monophyletic, species cluster in species-groups as proposed by morphology, and within each of the species groups the relations among the species are well supported. In addition, *COI* supports the hypothesis of the sister-group relationship between “castaneum” and “confusum” groups.

We also analyzed the effects of several weighting schemes; all produced the same tree topology, so we proceeded with the analyses by using simultaneously two weighting methodologies, substitution weighting 2 : 1 for T_v/T_s and character weighting 2 : 5 : 1 for the first vs. second vs. third codon positions, being one of the most commonly used weighting schemes. Ignoring transitions and substitutions on third codon positions resulted in low phylogenetic resolution, indicating that such an approach results in a loss of information.

16S rDNA

The trees obtained by all three methods for 16S rDNA gene revealed fairly similar topologies (not shown). The major uncertainty concerned the position of *T. brevicornis*, which was separated as a species group or joined to the “confusum” clade. 16S is phylogenetically most suitable for resolving relationships among taxa exhibiting intermediate levels of *COI* divergence (~5–15%) (Funk et al., 1995; Funk, 1999). However, *COI* divergence between species of *Tribolium* was 12–19%, and the 16S marker alone turned out to be insufficient to resolve unequivocally the phylogeny of the genus.

Combined *COI* and 16S rDNA

Since separate analysis of mitochondrial *COI* and 16S genes could not completely resolve the phylogenetic relationships among *Tribolium* species under study, further improvement was obtained by the analysis of combined 642 bp long *COI* fragment plus 448–452 bp long 16S fragment. The ILD test (Farris et al., 1994) rendered a *P* value of 0.168, which is nonsignificant, sug-




Fig. 1. Phylogenetic relationships among eight examined *Tribolium* species proposed on the basis of combined analysis of *COI* and 16S data sets. Acronyms “p” and “h” in the names of clones denote population and haplotype numbers, respectively. (a) The strict consensus MP tree (length 1178, CI = 0.610, RI = 0.852); confidence levels of tree bifurcation were assessed by 1000 bootstrap replicates. Bootstrap values are indicated above the nodes, while the partitioned Bremer support values are given below the nodes (*COI*/16S). Two pairs of sibling species (Hinton, 1948; Halstead, 1969) are indicated with asterisks, and proposed species groups with their geographic regions of origin (Hinton, 1948) are indicated with vertical bars. (b) Majority-rule consensus tree resulting from Bayesian MCMC analysis. Numbers on the nodes are posterior probability values. Branch lengths are proportional to number of changes.

gesting that the two data sets are combinable. Partitioned Bremer support values (Fig. 1a) indicate that the *COI* partition makes a slightly higher contribution to the overall support for the nodes than does the 16S partition.

The trees obtained in all three MP (Fig. 1a), ML (not shown) and Bayesian (Fig. 1b) analyses of combined sequences clearly showed monophyly of the genus *Tribolium* and also of each of the species. “Castaneum” and “confusum” clades are well supported, and within these two groups the relationships between the species as well as the two sibling species pairs (*T. freemani* – *T. castaneum* and *T. madens* – *T. audax*) are clearly resolved. Most nodes have high confidence values, except for the position of *T. brevicornis*. This species is positioned as a sister taxon to all other *Tribolium* species with a moderate support in MP and ML trees, while in the Bayesian tree it appears as a sister taxon to “castaneum” clade (though with low support).

DISCUSSION

Despite its great economical importance, the Tenebrionid genus *Tribolium* has not been extensively studied so far on the level of informational macromolecules, except for highly repetitive DNA traits of the several stored-product pest species (Juan et al., 1993; Ugarković et al., 1996; Mravinac et al., 2004, 2005). The reason for this neglect most probably resides in fact that the great majority of *Tribolium* species are free-living, many of them are very rare and/or inhabit remote and not easily reachable areas, and, to our knowledge, very few people might have (or, have had) the access to these species. This is especially so for the species belonging to “myrmecophilum” and “alcine” groups (restricted to Australia and Madagascar, respectively), but also for the “brevicornis” group species with the exception of *T. brevicornis*. Our original intention was to collect and include in our phylogenetic analysis the representatives of all *Tribolium* species-groups, but, unfortunately, this task turned out to be impossible. However, despite of this initial failure, we

decided to conduct this study in order to establish a preliminary phylogeny that could in turn be expanded and completed by the addition of the missing taxa.

In our study we included eight *Tribolium* species belonging to three species groups – “confusum”, “castaneum” and “brevicornis”. In order to resolve the phylogenetic relationships of these species we analyzed the most conserved part of the COI gene and examined part of 16S rDNA which evolves at least twice as slow as COI and is suitable for analysis of divergent species within a genus, differentiating among genera and families (Funk, 1999). Separate analyses of these two mitochondrial genes resulted in similar classification of *Tribolium* species, but, taken independently, they were not sufficient to resolve the phylogenetic relationships of the eight species under study. The phylogenetic relationships among examined *Tribolium* species were therefore analyzed using the combined segments of COI and 16S rDNA genes. Such an analysis showed monophyly of the genus *Tribolium* and clearly resolved species in three groups (Fig. 1), mostly in the same manner as they have been clustered according to the examination of morphological characters (Hinton, 1948). In the absence of the “myrmecophilum” and “alcine”-group species it is obviously not possible to confirm the sister-group relationships which would involve these species-groups, but the results of MP and ML suggest, with a moderate support, that the “brevicornis” group is probably the sister group to the other *Tribolium* species clade. However, since the Bayesian analysis suggested a slight possibility of a closer relationship of *T. brevicornis* with the “castaneum” group, we regard that the exact position of *T. brevicornis* could not be unequivocally resolved. We believe that the addition of species belonging to “brevicornis” group would greatly enhance its resolution.

Phylogenetic analysis of thirty-seven coleopteran species based on the mt COI gene revealed very long branches for species belonging to the families Scarabaeidae and Tenebrionidae, indicating high divergence among members of these families (Howland & Hewitt, 1995). Although the present study involved relatively conserved segments of COI and 16S rDNA mitochondrial genes, the sequence divergence was surprisingly high among examined *Tribolium* species, in a range from 12 to 19% for COI and from 7 to 18% for 16S. Low intraspecific variability is observed for both tested genes in all analyzed *Tribolium* species, including laboratory and wild populations, which can be almost exclusively found in storehouses (Wool, 1982).

Phylogenetic analysis of the collembolan genus *Orchesella* (Frati et al., 2000) and coleopteran genera *Neochlamisus*, *Ophraella*, *Timarcha* and *Chrysolina* (Funk et al., 1995; Funk, 1999; Garin et al., 1999; Gómez-Zurita et al., 2000) revealed intrageneric divergences in a high range from approx. 0.7% to 21% for COI and from 0.2% to 12% for 16S rDNA. In contrast to the insect species mentioned above, observed divergences are dramatically pronounced even among very close relatives within *Tribolium* species: siblings *T. madens* – *T. audax* and *T. freemani* – *T. castaneum*. For these species pairs, observed divergence is 14% and 15% for COI, and 7% and 13.5% for 16S rDNA, respectively, being within the level of divergence that has been usually observed between species groups in coleopteran genera (Funk et al., 1999). In species of the genus *Pissodes* (Coleoptera), which can hybridize producing fertile hybrids, the divergence of COI ranges from 0.5% to 7.5% (Langor & Sperling, 1997). Although COI in *Tribolium* showed higher sequence divergence than observed in other coleopteran species (e.g., Funk et al., 1995; Funk, 1999; Garin et al., 1999; Gómez-Zurita et al., 2000), the maximum divergence of COI at the amino acid level within *Tribolium* amounts to 7%, being within the divergence of 11% predicted for species within cole-

opteran genera. This is in agreement with the notion that COI is the most slowly evolving among mitochondrial protein coding genes at the amino acid level (Funk, 1999).

Based on the facts presented here, we assume that COI and 16S rRNA genes are not sufficient to completely resolve the phylogeny of the genus *Tribolium*. Future work on this genus should therefore include additional molecular markers, preferably some of the nuclear genes or genomic regions that proved useful for investigations of sub-familial interrelationships of insects, such as ribosomal RNA genes, elongation factors or ITS regions (e.g. Danforth et al., 2005; Gómez-Zurita & Galián, 2005; Gómez-Zurita et al., 2005; Becerra, 2004; Whitfield et al., 2002; Rokas et al., 2002; Cryan et al., 2001; Clark et al., 2001; Krzywinski et al., 2001; Mardulyn & Whitfield, 1999).

In conclusion, our study gives moderate support to the topology of the phylogenetic tree based on morphological traits proposed by Hinton (1948). We strongly hope that this preliminary analysis of phylogenetic relationships within this genus will serve as a stimulus for the *Tribolium*-oriented community to initiate a more thorough taxon sampling of wild-living species of *Tribolium* in order to fill the currently existing gaps and to reveal the “true” phylogeny of this important beetle-genus.

ACKNOWLEDGMENTS. We thank C. Juan and J. Pons for useful comments on the previous version of the manuscript. Critical reviews of two anonymous referees significantly improved our work. This study was supported by the Research Fund of Republic of Croatia, project No. P00981006.

REFERENCES

- BAKER R.H. & DeSALLE R. 1997: Multiple sources of character information and the phylogeny of Hawaiian drosophilids. *Syst. Biol.* **46**: 654–673.
- BECCERRA J.X. 2004: Molecular systematics of Blepharida beetles (Chrysomelidae: Alticinae) and relatives. *Mol. Phyl. Evol.* **30**: 107–117.
- BEEMAN R.W., STUART J.J., HAAS M.S. & FRIESEN K.S. 1996: Chromosome extraction and revision of linkage group 2 in *Tribolium castaneum*. *J. Hered.* **87**: 224–232.
- BENNETT R.L., BROWN S.J. & DENELL R.E. 1999: Molecular and genetic analysis of the *Tribolium* ultrabithorax ortholog, ultratorax. *Dev. Genes Evol.* **209**: 608–619.
- BENSASSON D., ZHANG D.-X., HARTL D.L. & HEWITT G.M. 2001: Mitochondrial pseudogenes: evolution’s misplaced witnesses. *Trends Ecol. Evol.* **16**: 314–321.
- BROWN S., DeCAMILIS M., GONZALES-CHARNECO K., DENELL M., BEEMAN R. & NIE W. 2000: Implications of the *Tribolium* deformed mutant phenotype for the evolution of Hox gene function. *Proc. Natl. Acad. Sci. USA* **97**: 4510–4514.
- BROWN S.J., DENELL R.E. & BEEMAN R.W. 2003: Beetling around the genome. *Genet. Res.* **82**: 155–161.
- CLARK T.L., MEINKE L.J. & FOSTER J.E. 2001: Molecular phylogeny of Diabrotica beetles (Coleoptera: Chrysomelidae) inferred from analysis of combined mitochondrial and nuclear DNA sequences. *Insect Mol. Biol.* **10**: 303–314.
- CRYAN J.R., LIEBHERR J.K., FETZNER J.W. & WHITING M.F. 2001: Evaluation of relationships within the endemic Hawaiian Platynini (Coleoptera: Carabidae) based on molecular and morphological evidence. *Mol. Phyl. Evol.* **21**: 72–85.
- DANFORTH B.N., LIN C.P. & FANG J. 2005: How do insect nuclear ribosomal genes compare to protein-coding genes in phylogenetic utility and nucleotide substitution patterns? *Syst. Entomol.* **30**: 549–562.
- DE BRUIJN M.H.L. 1983: *Drosophila melanogaster* mitochondrial DNA, a novel gene organisation and genetic code. *Nature* **304**: 234–241.

- DE RIJK P., VAN DE PEER Y. & DE WACHTER R. 1996: Database on the structure of large ribosomal subunit RNA. *Nucleic Acids Res.* **24**: 92–97.
- FARRIS J.S., KALLERSJO M., KLUGE A.G. & BULT C. 1994: Testing significance of incongruence. *Cladistics* **10**: 315–319.
- FELSENSTEIN J. 1985: Confidence limits on phylogenies: An approach using the bootstrap. *Evolution* **39**: 783–791.
- FRATI F., DELL'AMPIO E., CASASANTA S., CARAPPELLI A. & FANCIULLI P.P. 2000: Large amounts of genetic divergence among Italian species of the genus *Orchesella* (Insecta, Colembola) and the relationships of two new species. *Mol. Phyl. Evol.* **17**: 456–461.
- FUNK D.J. 1999: Molecular systematics of cytochrome oxidase I and 16S from *Neochlamisus* leaf beetles and the importance of sampling. *Mol. Biol. Evol.* **16**: 67–82.
- FUNK D.J., FUTUYUMA D.J., ORTI G. & MEYER A. 1995: Mitochondrial DNA sequences and multiple data sets: a phylogenetic study of phytophagous beetles (Chrysomelidae: Ophraella). *Mol. Biol. Evol.* **12**: 627–640.
- GARIN C.F., JUAN C. & PETITPIERRE E. 1999: Mitochondrial DNA phylogeny and the evolution of host-plant use in palearctic *Chrysolina* (Coleoptera, Chrysomelidae) leaf beetles. *J. Mol. Evol.* **48**: 435–444.
- GÓMEZ-ZURITA J., JUAN C. & PETITPIERRE E. 2000: The evolutionary history of the genus *Timarcha* (Coleoptera, Chrysomelidae) inferred from mitochondrial COII gene and partial 16S rDNA sequences. *Mol. Phyl. Evol.* **14**: 304–317.
- GÓMEZ-ZURITA J. & GALIÁN J. 2005: Current knowledge on genes and genomes of phytophagous beetles (Coleoptera: Chrysomeloidea, Curculionoidea): a review. *Eur. J. Entomol.* **102**: 577–597.
- GÓMEZ-ZURITA J.I., JOLIVET P. & VOGLER A.P. 2005: Molecular systematics of Eumolpinae and the relationships with Spilopyrinae (Coleoptera, Chrysomelidae). *Mol. Phyl. Evol.* **34**: 584–600.
- GUTELL R.R. & FOX G.E. 1988: A compilation of large subunit RNA sequences presented in a structural format. *Nucleic Acids Res. (Suppl.)* **16**: r175–r269.
- GUTELL R.R., LARSEN N. & WOESE C.R. 1994: Lessons from an evolving rRNA: 16S and 23S rRNA structures from a comparative perspective. *Microbiol. Rev.* **58**: 10–26.
- HALSTEAD D.G.H. 1969: A new species of *Tribolium* from North America previously confused with *Tribolium madens* (Chapr.) (Coleoptera: Tenebrionidae). *J. Stored Prod. Res.* **3**: 269–272.
- HALL T.A. 1999: BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucl. Acids. Symp. Ser.* **41**: 95–98.
- HINTON H.E. 1948: A synopsis of the genus *Tribolium* MacLeay with some remarks on the evolution of its species groups. *Bull. Entomol. Res.* **39**: 13–55.
- HOWARD R.D. 1987: Chemosystematic studies of the Triboliini (Coleoptera: Tenebrionidae): phylogenetic inferences from the defensive chemicals of eight *Tribolium* spp., *Palorus ratzeburgii* (Wissmann), and *Latheticus oryzae* Waterhouse. *Ann. Entomol. Soc. Am.* **80**: 398–405.
- HOWLAND D.E. & HEWITT G.M. 1995: Phylogeny of the Coleoptera based on mitochondrial cytochrome oxidase I sequence data. *Insect Mol. Biol.* **4**: 203–215.
- HUELSENBECK J.P. & RONQUIST F.R. 2001: MrBayes: Bayesian inference of phylogeny. *Bioinformatics* **17**: 754–755.
- JUAN C., VÁZQUEZ P., RUBIO J.M., PETITPIERRE E. & HEWITT G.M. 1993: Presence of highly repetitive DNA sequences in *Tribolium* flour-beetles. *Heredity* **70**: 1–8.
- JUAN C., OROMÍ P. & HEWITT G.M. 1995: Mitochondrial DNA phylogeny and sequential colonisation of Canary Islands by darkling beetles of the genus *Pimelia*. *Proc. R. Soc. Lond. (B)* **261**: 173–180.
- JUAN C., OROMÍ P. & HEWITT G.M. 1996: Phylogeny of the genus Hegeter (Tenebrionidae, Coleoptera) and its colonisation of the Canary Islands deduced from cytochrome oxidase I mitochondrial DNA sequences. *Heredity* **76**: 392–403.
- KRZYWINSKI J., WILKERSON R.C. & BESANSKY N.J. 2001: Evolution of mitochondrial and ribosomal gene sequences in Anophelinae (Diptera: Culicidae): implications for phylogeny reconstruction. *Mol. Phyl. Evol.* **18**: 479–487.
- KUMAR S., TAMURA K., JAKOBSEN I.B. & NEI M. 2001: *MEGA2: Molecular Evolutionary Genetics Analysis Software*. Arizona State University, Tempe, Arizona, USA.
- LANGOR D.W. & SPERLING F.A.H. 1997: Mitochondrial DNA sequence divergence in weevils of the *Pissodes strobi* species complex (Coleoptera, Curculionidae). *Insect Mol. Biol.* **6**: 255–265.
- LIU H. & BECKENBACH A.T. 1992: Evolution of the mitochondrial cytochrome oxidase II gene among 10 orders of insect. *Mol. Phyl. Evol.* **1**: 41–52.
- LUNT D.H., ZHANG D.-X., SZYMURA J.M. & HEWITT G.M. 1996: The insect cytochrome oxidase I gene: evolutionary patterns and conserved primers for phylogenetic studies. *Insect Mol. Biol.* **5**: 153–165.
- MARDULYN P. & WHITFIELD J.B. 1999: Phylogenetic signal in the COI, 16S, and 28S genes for inferring relationships among genera of Microgastrinae (Hymenoptera; Braconidae): evidence of a high diversification rate in this group of parasitoids. *Mol. Phyl. Evol.* **12**: 282–294.
- MEŠTROVIĆ N., MRAVINAC B., JUAN C., UGARKOVIĆ Đ. & PLOHL M. 2000: Comparative study of satellite sequences and phylogeny of five species from the genus *Palorus* (Insecta, Coleoptera). *Genome* **43**: 776–785.
- MRAVINAC B., PLOHL M. & UGARKOVIĆ Đ. 2004: Conserved patterns in the evolution of *Tribolium* satellite DNAs. *Gene* **332**: 169–177.
- MRAVINAC B., UGARKOVIĆ Đ., FRANJEVIĆ D. & PLOHL M. 2005: Long inversely oriented subunits form a complex monomer of *Tribolium brevicornis* satellite DNA. *J. Mol. Evol.* **60**: 513–525.
- NAGARAJU J. & GOLDSMITH M. R. 2002: Silkworm genomics – progress and prospects. *Current Sci.* **83**: 415–425.
- PONS J., BRUVO B., PETITPIERRE E., PLOHL M., UGARKOVIĆ Đ. & JUAN C. 2004: Complex structural features of satellite DNA sequences in the genus *Pimelia* (Coleoptera, Tenebrionidae): random differential amplification from a common “satellite DNA library”. *Heredity* **92**: 418–427.
- POSADA D. & CRANDALL K.A. 1998: MODELTEST: testing the model of DNA substitution. *Bioinformatics* **14**: 817–818.
- ROKAS A., NYLANDER J.A.A., RONQUIST F. & STONE G.N. 2002: A maximum-likelihood analysis of eight phylogenetic markers in gallwasps (Hymenoptera: Cynipidae): implications for insect phylogenetic studies. *Mol. Phyl. Evol.* **22**: 206–219.
- SCHMID K.J. & TAUTZ D. 1999: A comparison of homologous developmental genes from *Drosophila* and *Tribolium* reveals major differences in length and trinucleotide repeat content. *J. Mol. Evol.* **49**: 558–566.
- SIMMONS M.P. & OCHOTERENA H. 2000: Gaps as characters in sequence-based phylogenetic analyses. *Syst. Biol.* **49**: 369–381.
- SIMMONS M.P., OCHOTERENA H. & CARR T.G. 2001: Incorporation, relative homoplasy, and the effect of gap characters in

- sequence-based phylogenetic analyses. *Syst. Biol.* **50**: 454–462.
- SIMON C., FRATI F., BECKENBACH A., CRESPI B., LIU H. & FLOOK P. 1994: Evolution, weighting and phylogenetic utility of mitochondrial gene sequences and a compilation of conserved PCR primers. *Ann. Entomol. Soc. Am.* **87**: 651–701.
- SMITH S.G. 1952: The cytology of some tenebrionid beetles (Coleoptera). *J. Morphol.* **91**: 325–364.
- SOKOLOFF A. 1972: *The Biology of Tribolium, with Special Emphasis on Genetic Aspects*. Oxford University Press, Oxford, 300 pp.
- SORENSEN M.D. 1999: *TreeRot, Version 2. Computer software and documentation*. Boston University, MA.
- STEINER J.J., POKLEMB A.C.J., FJELLSTROM R.G. & ELLIOTT L.F. 1995: A rapid one-tube genomic DNA extraction process for PCR and RAPD analyses. *Nucl. Acids Res.* **23**: 2569–2570.
- SWOFFORD D.L. 1998: *PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods)*. Version 4. Sinauer Associates, Sunderland, Massachusetts.
- THOMPSON J.D., GIBSON T.J., PLEWNIAK F., JEANMOUGIN F. & HIGGINS D.G. 1997: The Clustal-X-Windows interface: Flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucl. Acids Res.* **25**: 4876–4882.
- UGARKOVIĆ Đ., PODNAR M. & PLOHL M. 1996: Satellite DNA of the red flour beetle-Tribolium castaneum and comparative study of satellites from the genus Tribolium. *Mol. Biol. Evol.* **13**: 1059–1066.
- WADE M.J. & JOHNSON N.A. 1994: Reproductive isolation between two species of flour beetles, Tribolium castaneum and T. freemani: variation within and among geographical populations of T. castaneum. *Heredity* **72**: 155–162.
- WHITFIELD J.B., MARDULYN P., AUSTIN A.D. & DOWTON M. 2002: Phylogenetic relationships among microgastrine braconid wasp genera based on data from the 16S, COI and 28S genes and morphology. *Syst. Entomol.* **27**: 337–359.
- WOOL D. 1982: Critical examination of postulated cladistic relationships among species of flour beetles (Tribolium, Tenebrionidae, Coleoptera). *Biochem. Genet.* **20**: 333–349.
- YOUNG N. & HEALY J. 2002: *GapCoder – A computer program for including indels in phylogenetic analysis*. <http://www.tufts.edu/vet/richlab/young/GapCoder/>
- ZHANG D-X. & HEWITT G.M. 1997: Assessment of the universality and utility of a set of conserved mitochondrial COI primers in insects. *Insect Mol. Biol.* **6**: 143–150.

Received July 11, 2005; revised and accepted December 23, 2005