

ERRATUM

HADDRILL P.R., SHUKER D.M., MAYES S. & MAJERUS M.E.N. 2007: Temporal effects of multiple mating on components of fitness in the two-spot ladybird, *Adalia bipunctata* (Coleoptera: Coccinellidae). *Eur. J. Entomol.* **104**(3): 393–398.

Some values in Table 1 in the above paper were unfortunately reversed. The correct version of Table 1 is reprinted below.

TABLE 1. Descriptive statistics from the experiment with respect to treatment. Not all females survived to complete the experiment. The mean number of days a female spent in the experiment is therefore given here. The total number of females surviving the 25 days of the experiment for each treatment is also given. For further details see main text. Arithmetic means and standard errors are presented except in the case of proportion data, where binomial means and standard errors are presented.

Treatment	Mean egg production	Mean no. of hatched eggs	Mean proportion of eggs hatched	Mean days in the experiment	No. of females surviving	Mean copulation duration	Mean no. of cycles
SM (n = 9)	410.67 ± 107.12	245.22 ± 75.46	0.60 ± 0.01	20.00 ± 2.96	6	184.11 ± 23.10	1.78 ± 0.28
RM (n = 9)	554.56 ± 95.14	290.67 ± 71.24	0.52 ± 0.01	22.89 ± 1.51	7	223.86 ± 9.70	2.31 ± 0.09
MM (n = 9)	441.78 ± 85.28	317.22 ± 64.87	0.72 ± 0.01	20.11 ± 2.75	6	222.41 ± 9.57	2.20 ± 0.09