



Locations of Rab, Allatotropin, Prothoracicotropic hormone and Period in the larval brain, corpus allatum and frontal ganglion of *Bombyx mori*

ASUKA MATSUI^{1,*}, MAKOTO TOKUSHIGE^{1,*}, AKIRA MIZOGUCHI², KENGO KANAMARU¹, KATSUHIKO SAKAMOTO³, YUICHI UNO⁴ and TOMOHIDE UNO^{1,**}

¹ Department of Applied Chemistry in Bioscience Biological Chemistry, Faculty of Agriculture, Kobe University, Nada-ku, Kobe, Hyogo 657-8501, Japan; e-mails: 213a402a@stu.kobe-u.ac.jp, 202a428a@stu.kobe-u.ac.jp, kng@kobe-u.ac.jp, unotom@kobe-u.ac.jp

² Division of Liberal Arts and Sciences, Aichi Gakuin University, Nissin, Aichi 470-0195, Japan; e-mail: amizo@dpc.agu.ac.jp

³ Graduate School of Agricultural Science, Faculty of Agriculture, Kobe University, Nada-ku, Kobe, Hyogo 657-8501, Japan; e-mail: ksakamoto@kobe-u.ac.jp

⁴ Department of Plant Resource Science, Faculty of Agriculture, Kobe University, Nada-ku, Kobe, Hyogo 657-8501, Japan; e-mail: yuno@kobe-u.ac.jp

Key words. Lepidoptera, Bombycidae, immunohistochemistry, silkworm, frontal ganglion, insect neurohormone, PTTH

Abstract. Rab, a low molecular weight GTP-binding protein, regulates the transmission of intracellular proteins. Insect neuro-peptides that are directly involved in growth, development, reproduction, homeostasis, courtship, feeding, circadian rhythm and many other physiological processes are synthesized in neurons and ganglia in the brain and secreted by specific neurosecretory cells in tissues such as the corpus allatum and frontal ganglion. To clarify the relationship between Rabs, two neuropeptides, Allatotropin (AT) and Prothoracicotropic hormone (PTTH) and the circadian clock protein, Period (PER), were used to determine the locations of nine Rabs (Rab1, Rab3, Rab6, Rab7, Rab11, Rab14, Rab19, Rab21 and RabX4). Rab6-, Rab11-, Rab14- and Rab21-immunohistochemical reactivities (IRs) partially overlapped AT-IR in the brain in *B. mori*. Rab3-, Rab6-, Rab11-, Rab14-, and Rab21-IRs overlapped AT-IR in the frontal ganglion. Of the seven Rabs, only Rab11-IR overlapped PTTH-IR in the brain. Rab1-, Rab3-, Rab11-, Rab14-, Rab19-, Rab21-, RabX4- and NUF (nuclear fallout, an effector of Rab11)-IRs, overlapped PER-IR in the brain. Therefore, Rab may regulate the exocytosis of PTTH, AT and a protein associated with the circadian rhythm.

INTRODUCTION

Small GTPases function as regulators of various cellular events by cycling between two nucleotide-states: a GDP-bound inactive form and a GTP-bound active form (Barr & Lambright, 2010; Müller & Goody, 2018). The largest family of small GTPases is Rab, which in mammals includes around 60 proteins (Stenmark, 2009; Pfeffer, 2017). Rab small GTPases regulate intracellular vesicular transmission (Barr, 2013; Li & Marlin, 2015; Pfeffer, 2017; Jin et al., 2021). Each Rab specifically targets a distinct component of a membrane. For example, Rab11 regulates recycling and secretion of membrane proteins between the plasma membrane and endosomes, Rab6 regulates transport in the Golgi apparatus, Rab7 regulates membrane fusion between vesicles and lysosomes and is involved in autophagy, and Rab11 regulates the traffic and transport of recycling endosomes. Rab14 is located in Golgi/TGN

and early endosomes. Rab21 is involved in endocytosis and early endosomal transport and Rab1 regulates transport from the endoplasmic reticulum to the Golgi apparatus (Plutner et al., 1991; Ullrich et al., 1996; Junutula et al., 2004; Simpson et al., 2004; Grigoriev et al., 2007; Wang et al., 2011). These Rabs are also involved in various phenomena in insects, including differentiation of various organs, fertilization, development, dormancy and secretion of neurotransmitters and neuropeptides. (Lighthouse et al., 2008; Uytterhoeven et al., 2011; Ye et al., 2012; Garg & Wu, 2014; Fujita et al., 2017; Elmogy et al., 2018; Caviglia et al., 2019).

Genome analyses indicate that insects have over 40 neuropeptides, for example, there are 42 in *Drosophila* (Nassel & Winther, 2010), 67 in *Apis mellifera* (Boerjan et al., 2010), 36 in *Pteromalus puparum* (Xu et al., 2020) and 67 in *Periplaneta Americana* (Zeng et al., 2021). These neuro-

* These authors contributed equally to this work.

** Corresponding author; e-mail: unotom@kobe-u.ac.jp

peptides are synthesized in specific neurons, mainly in the brain, and are transported along axons to terminal neurosecretory cells in structures such as the corpus allatum (CA) and frontal ganglion (FG). They are then secreted into the haemolymph or their information is transmitted to different neurons (Nassel, 2002; Heuer et al., 2012; Bednar et al., 2017; Schoofs et al., 2017).

The neuropeptide Allatotropin (AT), was isolated based on its activity in stimulating juvenile hormone synthesis in the CA of the lepidopteran *Manduca sexta* (Kataoka et al., 1989). It is expressed in the brain and frontal ganglion (FG), is multifunctional, acting in different insects as a myoregulator and a cardioaccelerator (Duve et al., 1999; Koladich et al., 2002; Villalobos-Sambucaro et al., 2015). AT is synthesized in a number of neurons in the central nervous system and two neurons in the FG (Nagata et al., 2012; Bednar et al., 2017); however, the molecular mechanisms controlling its transport and secretion are unknown.

Prothoracicotropic hormone (PTTH) stimulates the endocrine prothoracic gland to secrete the metamorphic hormone, ecdysone, into the haemolymph. PTTH plays a central role in the endocrine network, controlling insect growth, moulting, metamorphosis and diapause (Goltzené et al., 1992; Nagata et al., 2005). PTTH is released from the CA, which is an organ specific to insects, which together with associated nerves secretes neuropeptides into the haemolymph (Mizoguchi et al., 1987; O'Brien et al., 1988).

Clock cells in the dorsolateral protocerebrum (DL) in the brain express clock genes, such as *Period* (*Per*), *Timeless* and *Clock*, which secrete the neuropeptide Pigment dispersing factor (PDF), which regulates the biological clock of insects (Sehadová et al., 2004; Hardin, 2005; Iwai et al., 2008; Inami et al., 2022). PTTH was identified in several lepidopterans as a product of two pairs of brain neurons that are distinct from those expressing *Per* and *Timeless* (Sauman & Reppert, 1996; Wise et al., 2002). Clock neurons can regulate the synthesis and rhythmic release of PTTH into the haemolymph (Vafopoulou & Steel, 2014).

Immunohistochemical analyses using antibodies against several Rab proteins have revealed Rab co-localisation with Bombyxin, an insulin-like peptide, in neurons in the brain and CA of the silkworm (Uno et al., 2013, 2021). However, the localization of Rab using neuropeptides other than Bombyxin is rare. Therefore, to clarify the role of Rab in neuropeptide secretion, an immunohistochemical analysis was carried out using antibodies against multiple Rabs and the neuropeptides, PTTH and AT, and clock protein, PER.

MATERIALS AND METHODS

Materials and insect cultures

Oligonucleotides were purchased from Invitrogen (Tokyo, Japan). The other chemicals used were of the purest grade commercially available. We used hybrid (Kinshu × Showa or Daizo, p50 strains) *B. mori* and 5 day-old fourth-instar larvae in all the experiments.

Antiserum production

Antisera against *B. mori* Rab1, 3, 6, 7, 11, 14, 21, X4, and PTTH, PER and AT were obtained as previously described (Uno et al., 2010, 2013, 2016, 2017, 2021). The specificity of the antibodies is shown in Table 1. A fragment of cDNA containing the partial coding sequence of *B. mori* Nuf (*Nuclear fallout*) and Rab19 was generated using reverse transcriptase-PCR (RT-PCR), subcloned and sequenced. This fragment was inserted between the *Bam*HI and *Eco*RI sites of pGEX6P-2 (GE Healthcare UK Ltd, UK) and expressed in *E. coli* (BL21 strain). NUF and Rab19 GST-protein was purified using a glutathione S-Sepharose column as previously described (Uno et al., 2014). After digestion with PreScission protease, the His-tagged protein was purified using a His-accept column.

Antisera were generated in a rabbit by injecting it with a 1:1 (v/v) mixture of each purified *B. mori* protein (1 mg) and Freund's complete adjuvant. The rabbits received three booster injections at 2-weekly intervals. The sera were isolated and tested for the presence of anti-*B. mori* NUF and Rab19 antibodies using western blotting.

Western blotting and immunohistochemistry

Western blotting and immunohistochemistry were done as previously described (Uno et al., 2019). For western blotting, purified proteins (1 µg) were transferred to a PVDF membrane after electrophoresis. The membrane was placed in Blocking One solution for 60 min at room temperature [RT]) and then for 60 min at room temperature [RT]) with anti-NUF serum (1:2,000), as the primary antibody, in Tris-buffered saline (TBS; 50 mM Tris-HCl, 50 mM NaCl [pH 8.0]) containing Blocking One solution. The membrane was then washed (3×) with TBS (including 0.05% Tween-20 (v/v)), followed by incubation (60 min, room temperature [RT]) with the secondary antibodies: peroxidase-conjugated goat anti-rabbit IgG (both 1:2000). After the membrane was washed (3× with TBS plus Tween 20), proteins were detected using the peroxidase staining DAB kit (Nakalai Tesque Inc, Kyoto, Japan).

Insect heads were fixed (24 h at 4°C) in Bouin fluid (saturated picric acid, formalin and acetic acid at 15:5:1 by volume). Standard histochemical methods were used for dehydrating tissues and embedding in paraplast, then sectioning (9-µm-thick sections), followed by deparaffinization and rehydration. The insect heads were washed (at RT) in distilled water and in phosphate-buffered saline (PBS; 137 mM NaCl, 2.7 mM KCl, 10 mM Na₂HPO₄ and 2 mM KH₂PO₄, [pH 7.4]) containing 0.3% Triton-X100 (PBS-Tr), blocked (30 min at RT) with antibody dilution buffer (PBS-Tr containing 1.5% goat serum) and incubated (overnight at 4°C).

Table 1. The relationship between Rab and neuropeptides in terms of the specificity of the antibodies.

	Allatotropin		PTTH	Period	Bombyxin		Reference
	Brain	FG			Brain	CA	
Rab1	n.d.	n.d.	n.d.	+	+	n.d.	(1)
Rab3	n.d.	+	n.d.	+	+	+	(2)
Rab6	+	+	n.d.	n.d.	+	+	(2)
Rab7	n.d.	+	n.d.	–	+	+	(3)
Rab11	+	+	+	+	+	+	(4)
Rab14	+	+	n.d.	+	+	+	(1)
Rab19	n.d.	n.d.	n.d.	n.d.	–	–	(5)
Rab21	+	+	n.d.	+	+	+	(4)
RabX4	n.d.	n.d.	n.d.	+	+	+	(6)

n.d.: not detectable; –: not examined; +: colocalized. (1) Uno et al., 2013; (2) Uno et al., 2016; (3) Uno et al., 2010; (4) Uno et al., 2021; (5) This paper; (6) Uno et al., 2017.




Fig. 1. Co-localization of Rab1-, Rab3-, Rab6-, Rab7- and Rab11-IR with AT-IR in the brain of *B. mori*. Allatotropin (a, e, i, m and q) was visualized using CFTM555 (red fluorophore). Rab1 (b), Rab3 (f), Rab6 (j), Rab7 (n) and Rab11 (r) were visualized using CFTM488A (green fluorophore). Rab6- and Rab11-IRs partially colocalized in AT-IR neurons (k, l, x and y). Rab1-, Rab3- and Rab7-IRs did not overlap with AT-IR (c, d, g, h, o and p). Scale bar: 100 μ m.

in dilution buffer with each of the primary antibodies: anti-Rab rabbit serum (1:500), anti-NUF rabbit serum (1:500), anti-PER mouse IgG (1:500), anti-AT mouse IgG (1:500) and anti-PTTH mouse IgG (1:500). After rinsing (three times for 10 min at RT) with PBS-Tr, the sections were incubated (1.5 h at RT) with 7.5 μ g/ml of each of the secondary antibodies: donkey anti-rat IgG (H+L)-CF555 and goat anti-rabbit IgG (H+L)-CF488A. After washing in PBS-Tr, the sections were mounted in Aqua-Poly/Mount medium and examined using a BX50 microscope equipped with BX-FLA reflected light fluorescence and WIG and NIBA mirror/filter units. Three to five individuals were examined in each immunocytochemical experiment. The excitation wavelength and emission range in WIG mirror/filter unit were 520–550 nm and above 580 nm, respectively. The excitation wavelength and emission range in the NIBA mirror/filter unit were 470–490 nm and 515–550 nm, respectively.

In control experiments, the primary antibodies were replaced with pre-immune rabbit serum and no significant staining was observed above the background level. The bleed-through of fluorescence was checked by imaging singly stained samples using both filter/mirror units (WIG and NIBA). In both cases no significant staining was observed.

RESULTS

Location of immunoreactivity of Rabs and AT in the brain of *B. mori*

AT-IR was detected in many neural cells in the brain (Figs 1 and 2). This is consistent with previous findings reported for the silk moth (Bednar et al., 2017). Double-labelling showed that Rab6-, Rab11-, Rab14- and Rab21-IRs were partially colocalized with AT-IR (Figs 1 and 2,




Fig. 2. Co-localization of Rab14-, Rab19-, Rab21- and RabX4-IRs with AT-IR in the brain of *B. mori*. Allatotropin (a, e, i and m) was visualized using CFTM555 (red fluorophore). Rab14 (b), Rab19 (f), Rab21 (j) and RabX4 (n) were visualized using CFTM488A (green fluorophore). Rab14- and Rab21-IRs partially overlapped with AT-IR (c, d, k and l). Rab19- and RabX4-IRs did not overlap with AT-IR (g, h, o and p). Scale bar: 100 μ m.

arrow head). Rab1-, Rab3-, Rab7- and RabX4-IRs did not overlap with AT-IR. Rab6-IR and Rab21-IR were present in the ventral (Figs 1i–l, 2i–l), Rab11-IR in the dorso-lateral (Fig. 1q–y) and Rab14-IR in the central region of the brain (Fig. 2a–d).

Location of immunoreactivity of Rabs and AT in the FG of *B. mori*.

AT-IR was detected in two cells in the FG (Figs 3 and 4). This is consistent with previous findings reported for the silk moth (Bednar et al., 2017). Rab3-, Rab6-, Rab7-, Rab11-, Rab14- and Rab21-IRs overlapped with AT-IR in the FG (Figs 3 and 4, arrow head). Rab1-, Rab19- and RabX4-IRs did not overlap with AT-IR.

Location of immunoreactivity of Rabs and PTTH in the brain of *B. mori*

In Lepidoptera, PTTH is produced by two pairs of DL neurosecretory cells in the brain (Sauman & Reppert, 1996). PTTH-IR was detected in 1–2 neurons in the DL area (Fig. 5). Double-labelling showed that PTTH-IR colocalized with Rab11-IR in the brain (Fig. 5g–i, arrowhead) but not with other Rab-IR neurons in the brain (Fig.

5). Rab11-immunopositive neurons did not overlap with PTTH-IR in the CA (Fig. 5x).

Location of immunoreactivity of Rabs and PER in the brain of *B. mori*

Rab-IR cells did not overlap with PTTH-IR cells, but many Rab-IR cells were located close to PTTH-IR cells in the DL of the brain (Fig. 5). PER-IR cells were located close to the contralateral PTTH neurosecretory cells (Sauman & Reppert, 1996). Double-labelling showed that Rab1-, Rab3-, Rab11-, Rab14-, Rab19-, Rab21- and RabX4-IRs overlapped with PER-IR in the brain (Fig. 6).

Anti-*B. mori* NUF antibodies and location of NUF immunoreactivity

NUF is a Rab11 effector protein that is important in controlling the transmission of several intracellular cargos (Riggs et al., 2007; Calero-Cuenca & Sotillos, 2018).

As a control for binding specificity, the anti-NUF antibody was pre-incubated with an excess amount of antigen before immunological staining (Fig. 7d–f).

The bleed-through fluorescence was assessed by imaging the single-stained samples using both filter/mirror units (WIG and NIBA) (Fig. 7a–c). In both cases, no significant




Fig. 3. Co-localization of Rab1-, Rab3-, Rab6-, Rab7- and Rab11-IRs with AT-IR in the frontal ganglion of *B. mori*. Allatotropin (a, d, g, j and m) was visualized using CFTM555 (red fluorophore). Rab1 (b), Rab3 (e), Rab6 (h), Rab7 (k), and Rab11 (n) were visualized using CFTM488A (green fluorophore). Rab3-, Rab6-, Rab7- and Rab11-IRs overlapped with AT-IR (f, i, l and o). Rab1-IR did not overlap with AT-IR (c). Scale bar: 100 μ m.




Fig. 4. Co-localization of Rab14-, Rab19-, Rab21- and RabX4-IRs with AT-IR in the frontal ganglion of *B. mori*. Allatotropin (a, d, g, and j) was visualized using CFTM555 (red fluorophore). Rab14 (b), Rab19 (e), Rab21 (h), and RabX4 (k) were visualized using CFTM488A (green fluorophore). Rab14- and Rab21-IRs overlapped with AT-IR (c and i). Rab19- and RabX4-IRs did not overlap with AT-IR (f and l). Scale bar: 100 μ m.

staining was recorded (Fig. 7a, e). Antibodies produced against *B. mori* NUF specifically recognized the protein band corresponding to the position of purified partial *B. mori* NUF (Fig. 7p). Control experiments (the addition of the antigen and primary antibody together) did not result in staining above background levels (Fig. 7p).

Double-labelling showed that NUF-IR colocalized with Rab11-IR in neurons in the brain (Fig. 7g–i), but not with PTTH-IR neurons (Fig. 7j–l). NUF-IR also overlapped with PER-IR in the brain (Fig. 7m–o).

DISCUSSION

To clarify the relationship between Rabs, two neuropeptides, AT and PTTH, and the circadian clock protein, PER, the colocalization of nine Rabs (Rab1, Rab3, Rab6, Rab7, Rab11, Rab14, Rab19, Rab21 and RabX4) with these proteins was determined. Rab6-, Rab11-, Rab14- and Rab21-IRs partially overlapped with AT-IR in the brain. Rab3-, Rab6-, Rab7-, Rab11-, Rab14- and Rab21-IRs overlapped with AT-IR in the FG. Only Rab11-IR areas overlapped with PTTH-IR in the brain. Rab1-, Rab3-, Rab11-, Rab14-, Rab19-, Rab21- and RabX4-IRs and NUF-IR, an effector




Fig. 5. Co-localization of Rab1-, Rab3-, Rab11-, Rab14-, Rab19-, Rab21-, RabX4-IRs with PTTH-IR in the brain (a–u) and CA (v–x) of *B. mori*. PTTH (a, d, g, j, m, p, s and v) was visualized using CFTM555 (red fluorophore). Rab1 (b), Rab3 (e), Rab11 (h and w), Rab14 (k), Rab19 (n), Rab21 (q) and RabX4 (t) were visualized using CFTM488A (green fluorophore). Rab11-IR overlapped with PTTH-IR (i). Rab1-, Rab3-, Rab14-, Rab19-, Rab21- and RabX4-IRs did not overlap with AT-IR (c, f, l, o, r and u). Rab11-IR in the CA did not overlap with PTTH-IR (x). Scale bar: 100 μ m.




Fig. 6. Co-localization of Rab1-, Rab3-, Rab11-, Rab14-, Rab19-, Rab21- and RabX4-IRs with PER-IR in the brain of *B. mori*. PER (a, d, g, j, m, p and s) was visualized using CFTM555 (red fluorophore). Rab1 (b), Rab3 (e), Rab11 (h), Rab14 (k), Rab19 (n), Rab21 (q) and RabX4 (t) were visualized using CFTM488A (green fluorophore). Rab1-, Rab3-, Rab11-, Rab14-, Rab19-, Rab21-, and RabX4-IRs overlapped with PER-IR (c, c', f, f', i, i', l, l', o, o', r, r', u, and u'). Scale bar: 100 μ m.

of Rab11, overlapped with PER-IR in the brain. Table1 shows the relationship between Rabs and neuropeptides and the specificity of the antibodies.

AT-IR is present in many neurons in the central nervous system and in two neurons in the FG in *B. mori* (Nagata et al., 2012). Rab6-, Rab11-, Rab14- and Rab21-IRs partially overlapped AT-IR in the brain (Fig. 1 and 2). AT-IR was detected in two cells in the FG (Fig. 3 and 4). Rab3-, Rab6-,

Rab7-, Rab11-, Rab14- and Rab21-IRs overlapped AT-IR in the FG (Fig. 3 and 4). Rab6, Rab11, Rab14 and Rab21 in the brain may regulate the transport and secretion of AT.

Rab6 regulates transport in the Golgi apparatus and Rab11 the recycling and secretion of membrane proteins between the plasma membrane and endosomes. Rab6 and Rab11 may regulate the transport of AT in the brain and FG. Rab14 functions in Golgi/endosome transport process-




Fig. 7. Location and Immunoblot analysis of NUF in the brain of *B. mori*. Bleed-through fluorescence (a–c') and Control (d–f'). Single stained samples were imaged through both filters (a–c'). Nuf-IR was visualized using CF555 (red fluorophore, a) and CFTM 488A (green fluorophore, b). Control (d–f'); (antibody + antigen, used as a primary antibody). Nuf-IR was visualized using CFTM 488A (green fluorophore, e). Rab11-IR was visualized using CF555 (red fluorophore, a). In both cases, no significant staining was observed (c, c', f and f'). Rab11 (g), PTTH (j) and PER (m) were visualized using CFTM 555 (red fluorophore). NUF (h, k and n) was visualized using CFTM 488A (green fluorophore). NUF-IR colocalizes with Rab11-IR in neurons in the brain (i and i'), but not with PTTH-IR neurons in the brain (l and l'). Rab11-IR colocalized with PER-IR in neurons (o and o'). Scale bar: 100 μ m. Panel p, lanes 1–3 were loaded with His-NUF. Lane 2 was probed with pre-immune serum as a negative control. Lane 3 contains an antibody-positive antigen as a positive primary antibody control.

es and Rab21 in endocytosis and early endosomal transport (Plutner et al., 1991; Junutula et al., 2004; Simpson et al., 2004; Grigoriev et al., 2007; Wang et al., 2011). In addition, *Drosophila* Rab21 and Rab14 are associated with autophagy. *Drosophila* Rab21 promotes endosomal sorting of VAMP8 required for autophagosome-lysosome fusion and Rab14 is necessary for autophagosome growth, positioning and fusion (Jean et al., 2015; Mauvezin et al., 2016). Rab21 and Rab14 may be involved in autophagy-related degradation of AT in the brain and FG. Multiple Rabs may regulate transport and secretion of AT in the brain and FG. Further study is necessary to determine whether Rab effector-IR is present in AT-IR cells and whether the inhibition of Rabs by dsRNA interference inhibits AT secretion.

Of the Rabs, only Rab11 was present in cells synthesizing PTTH in the DL (Fig. 5). Rab11 is involved in membrane transmission of recycling endosomes. Rab11 proteins are involved in a great variety of cellular transmission pathways, such as, the regulation of cell polarity, ciliogenesis, integrin recycling, neuritogenesis, oogenesis, recep-

tor/adhesion protein recycling and cytokinesis (Simon & Prekeris, 2008; Jing & Prekeris, 2009; Eva et al., 2010; Knödler et al., 2010). As Rab11 is the only Rab present in PTTH-IR cells, PTTH secretion may be regulated by controlling vesicular membrane recycling. Rab11 may be involved downstream of the neurotransmitter serotonin because receptors for this neurotransmitter are also present in PTTH-IR cells (Wang et al., 2013).

NUF is a Rab11 effector protein that is important in controlling the transmission of several intracellular cargos (Riggs et al., 2007; Calero-Cuenca & Sotillos, 2018). NUF-IR cells did not colocalize with PTTH-IR (Fig. 7). Rab11 has multiple effectors (Dusty-like protein, Rip11) in addition to NUF (Nagaraj & Adler, 2012; Calero-Cuenca & Sotillos, 2018). These other effectors may be involved in PTTH secretion.

PTTH-IR was not present in Rab11-IR cells in the CA (Fig. 5). In the CA, another Rab may be involved in PTTH secretion. Alternatively, Rab may not be required at the site of PTTH secretion in the CA.

Except for Rab11-IR cells, Rab-IR was not present in PTTH-IR cells, but many Rab-IR cells in the DL were in close proximity to PTTH-IR cells (Fig. 5). Clock cells where PER is located are adjacent to PTTH synthesizing cells in DL. Rab1-, Rab3-, Rab11-, Rab14-, Rab19-, Rab21- and RabX4-IRs and NUF-IR overlapped with PER-IR in the brain (Figs 5 and 6).

Pigment dispersing factor (PDF), the only peptidergic hormone involved in circadian rhythms and a candidate transmitter of oscillations in behaviour and the clock proteins Timeless and Casein kinase, are located in these clock cells in DL (Iwai et al., 2008). In addition, synthesis and secretion of PTTH are regulated by circadian rhythms (Vafopoulou et al., 2007). Several Rabs in the clock cells in the brain may regulate the secretion and transport of PDF and circadian rhythm-related proteins.

To clarify the role of Rab in the secretion of neuropeptides, genetic methods, such as, direct introduction of dsRNA into the brain and gene editing need to be used to determine whether suppressing Rab reduces neuropeptide secretion or alters the biological clock. In addition, it is also necessary to determine whether effectors that specifically bind to Rab are located in the same cells in the brain.

CONFLICT OF INTERESTS. The authors declare that there are no conflict of interests.

ACKNOWLEDGEMENTS. We thank L. Benyon, from Edanz (<https://jp.edanz.com/ac>) for editing a draft of this manuscript. This work was supported by the Hyogo science and technology association.

REFERENCES

- BARR F.A. 2013: Review series: Rab GTPases and membrane identity: causal or inconsequential? — *J. Cell Biol.* **202**: 191–199.
- BARR F. & LAMBRIGHT D.G. 2010: Rab GEFs and GAPs. — *Curr. Opin. Cell Biol.* **22**: 461–470.
- BEDNAR B., ROLLER L., CIZMAR D., MITROVA D. & ZITNAN D. 2017: Developmental and sex-specific differences in expression of neuropeptides derived from allatotropin gene in the silkworm *Bombyx mori*. — *Cell Tissue Res.* **368**: 259–275.
- BOERJAN B., CARDOEN D., BOGAERTS A., LANDUYT B., SCHOOLS L. & VERLEYEN P. 2010: Mass spectrometric profiling of (neuro)peptides in the worker honeybee, *Apis mellifera*. — *Neuropharmacology* **58**: 248–258.
- CALERO-CUENCA F.J. & SOTILLOS S. 2018: Nuf and Rip11 requirement for polarity determinant recycling during *Drosophila* development. — *Small GTPases* **9**: 352–359.
- CAVIGLIA S., FLORES-BENITEZ D., LATTNER J., LUSCHNIG S. & BRANKATSCHK M. 2019: Rabs on the fly: Functions of Rab GTPases during development. — *Small GTPases* **10**: 89–98.
- DUVE H., EAST P.D. & THORPE A. 1999: Regulation of lepidopteran foregut movement by allatostatins and allatotropin from the frontal ganglion. — *J. Comp. Neurol.* **413**: 405–416.
- ELMOGY M., MOHAMED A.A., TUFAIL M., UNO T. & TAKEDA M. 2018: Molecular and functional characterization of the American cockroach, *Periplaneta americana*, Rab5: the first exopterygotan low molecular weight ovarian GTPase during oogenesis. — *Insect Sci.* **25**: 751–764.
- EVA R., DASSIE E., CASWELL P.T., DICK G., FRENCH-CONSTANT C., NORMAN J.C. & FAWCETT J.W. 2010: Rab11 and its effector Rab coupling protein contribute to the trafficking of beta 1 integrins during axon growth in adult dorsal root ganglion neurons and PC12 cells. — *J. Neurosci.* **30**: 11654–11669.
- FUJITA N., HUANG W., LIN T.H., GROULX J.F., JEAN S., NGUYEN J., KUCHITSU Y., KOYAMA-HONDA I., MIZUSHIMA N., FUKUDA M. ET AL. 2017: Genetic screen in *Drosophila* muscle identifies autophagy-mediated T-tubule remodeling and a Rab2 role in autophagy. — *Elife* **6**: e23367, 30 pp.
- GARG A. & WU L.P. 2014: *Drosophila* Rab14 mediates phagocytosis in the immune response to *Staphylococcus aureus*. — *Cell Microbiol.* **16**: 296–310.
- GOLTZENÉ F., HOLDER F., CHARLET M., MEISTER M. & OKA T. 1992: Immunocytochemical localization of *Bombyx*-PTTH-like molecules in neurosecretory cells of the brain of the migratory locust, *Locusta migratoria*. A comparison with neuropeptin and insulin-related peptide. — *Cell Tissue Res.* **269**: 133–140.
- GRIGORIEV I., SPLINTER D., KEIJZER N., WULF P.S., DEMMERS J., OHTSUKA T., MODESTI M., MALY I.V., GROSVELD F., HOOGENRAAD C.C. ET AL. 2007: Rab6 regulates transport and targeting of exocytotic carriers. — *Dev. Cell* **13**: 305–314.
- HARDIN P.E. 2005: The circadian timekeeping system of *Drosophila*. — *Curr. Biol.* **15**: R714–722.
- HEUER C.M., KOLLMANN M., BINZER M. & SCHACHTNER J. 2012: Neuropeptides in insect mushroom bodies. — *Arthropod Struct. Dev.* **41**: 199–226.
- INAMI S., SATO T. & SAKAI T. 2022: Circadian neuropeptide-expressing clock neurons as regulators of long-term memory: Molecular and cellular perspectives. — *Front. Mol. Neurosci.* **15**: 934222, 7 pp.
- IWAI S., THI DIEU TRANG L., SEHADOVA H. & TAKEDA M. 2008: Expression analyses of casein kinase 2alpha and casein kinase 2beta in the silkworm, *Bombyx mori*. — *Comp. Biochem. Physiol. (B, Biochem. Mol. Biol.)* **149**: 38–46.
- JEAN S., COX S., NASSARI S. & KIGER A.A. 2015: Starvation-induced MTMR13 and RAB21 activity regulates VAMP8 to promote autophagosome-lysosome fusion. — *EMBO Rep.* **16**: 297–311.
- JIN H., TANG Y., YANG L., PENG X., LI B., FAN Q., WEI S., YANG S., LI X., WU B. ET AL. 2021: Rab GTPases: Central coordinators of membrane trafficking in cancer. — *Front. Cell Dev. Biol.* **9**: 648384, 13 pp.
- JING J. & PREKERIS R. 2009: Polarized endocytic transport: the roles of Rab11 and Rab11-FIPs in regulating cell polarity. — *Histol. Histopathol.* **24**: 1171–1180.
- JUNUTULA J.R., DE MAZIERE A.M., PEDEN A.A., ERVIN K.E., ADVANI R.J., VAN DIJK S.M., KLUMPERMAN J. & SCHELLER R.H. 2004: Rab14 is involved in membrane trafficking between the Golgi complex and endosomes. — *Mol. Biol. Cell* **15**: 2218–2229.
- KATAOKA H., TOSCHI A., LI J.P., CARNEY R.L., SCHOOLEY D.A. & KRAMER S.J. 1989: Identification of an allatotropin from adult *Manduca sexta*. — *Science* **243**: 1481–1483.
- KNÖDLER A., FENG S., ZHANG J., ZHANG X., DAS A., PERÄNEN J. & GUO W. 2010: Coordination of Rab8 and Rab11 in primary ciliogenesis. — *Proc. Natl. Acad. Sci. U.S.A.* **107**: 6346–6351.
- KOLADICH P.M., CUSSON M., BENDENA W.G., TOBE S.S. & McNEIL J.N. 2002: Cardioacceleratory effects of *Manduca sexta* allatotropin in the true armyworm moth, *Pseudaletia unipuncta*. — *Peptides* **23**: 645–651.
- LI G. & MARLIN M.C. 2015: Rab family of GTPases. — *Methods Mol. Biol.* **2015**: 1298, 16 pp.
- LIGHTHOUSE D.V., BUSZCZAK M. & SPRADLING A.C. 2008: New components of the *Drosophila* fusome suggest it plays novel roles in signaling and transport. — *Dev. Biol.* **317**: 59–71.
- MAUVEZIN C., NEISCH A.L., AYALA C.I., KIM J., BELTRAME A., BRADEN C.R., GARDNER M.K., HAYS T.S. & NEUFELD T.P. 2016:

- Coordination of autophagosome-lysosome fusion and transport by a Klp98A-Rab14 complex in *Drosophila*. — *J. Cell Sci.* **129**: 971–982.
- MIZOGUCHI A., ISHIZAKI H., NAGASAWA H., KATAOKA H., ISOAGI A., TAMURA S., SUZUKI A., FUJINO M. & KITADA C. 1987: A monoclonal antibody against a synthetic fragment of bombyxin (4K-prothoracicotrophic hormone) from the silkworm, *Bombyx mori*: characterization and immunohistochemistry. — *Mol. Cell Endocrinol.* **51**: 227–235.
- MÜLLER M.P. & GOODY R.S. 2018: Molecular control of Rab activity by GEFs, GAPs and GDI. — *Small GTPases* **9**: 5–21.
- NAGARAJ R. & ADLER P.N. 2012: Dusky-like functions as a Rab11 effector for the deposition of cuticle during *Drosophila* bristle development. — *Development* **139**: 906–916.
- NAGATA S., KATAOKA H. & SUZUKI A. 2005: Silk moth neuropeptide hormones: prothoracicotrophic hormone and others. — *Ann. N. Y. Acad. Sci.* **1040**: 38–52.
- NAGATA S., MATSUMOTO S., MIZOGUCHI A. & NAGASAWA H. 2012: Identification of cDNAs encoding allatotropin and allatotropin-like peptides from the silkworm, *Bombyx mori*. — *Peptides* **34**: 98–105.
- NASSEL D.R. 2002: Neuropeptides in the nervous system of *Drosophila* and other insects: multiple roles as neuromodulators and neurohormones. — *Prog. Neurobiol.* **68**: 1–84.
- NASSEL D.R. & WINTNER A.M. 2010: *Drosophila* neuropeptides in regulation of physiology and behavior. — *Prog. Neurobiol.* **92**: 42–104.
- O'BRIEN M.A., KATAHIRA E.J., FLANAGAN T.R., ARNOLD L.W., HAUGHTON G. & BOLLENBACHER W.E. 1988: A monoclonal antibody to the insect prothoracicotrophic hormone. — *J. Neurosci.* **8**: 3247–3257.
- PFEFFER S.R. 2017: Rab GTPases: master regulators that establish the secretory and endocytic pathways. — *Mol. Biol. Cell* **28**: 712–715.
- PLUTNER H., COX A.D., PIND S., KHOSRAVI-FAR R., BOURNE J.R., SCHWANINGER R., DER C.J. & BALCH W.E. 1991: Rab1b regulates vesicular transport between the endoplasmic reticulum and successive Golgi compartments. — *J. Cell Biol.* **115**: 31–43.
- RIGGS B., FASULO B., ROYOU A., MISCHÉ S., CAO J., HAYS T.S. & SULLIVAN W. 2007: The concentration of Nuf, a Rab11 effector, at the microtubule-organizing center is cell cycle regulated, dynein-dependent, and coincides with furrow formation. — *Mol. Biol. Cell* **18**: 3313–3322.
- SAUMAN I. & REPERT S.M. 1996: Molecular characterization of prothoracicotrophic hormone (PTTH) from the giant silkworm *Antheraea pernyi*: developmental appearance of PTTH-expressing cells and relationship to circadian clock cells in central brain. — *Dev. Biol.* **178**: 418–429.
- SCHOOF L., DE LOOF A. & VAN HIEL M.B. 2017: Neuropeptides as regulators of behavior in insects. — *Annu. Rev. Entomol.* **62**: 35–52.
- SEHADOVÁ H., MARKOVA E.P., SEHNAL F. & TAKEDA M. 2004: Distribution of circadian clock-related proteins in the cephalic nervous system of the silkworm, *Bombyx mori*. — *J. Biol. Rhythms* **19**: 466–482.
- SIMON G.C. & PREKERIS R. 2008: Mechanisms regulating targeting of recycling endosomes to the cleavage furrow during cytokinesis. — *Biochem. Soc. Trans.* **36**: 391–394.
- SIMPSON J.C., GRIFFITHS G., WESSLING-RESNICK M., FRANSEN J.A., BENNETT H. & JONES A.T. 2004: A role for the small GTPase Rab21 in the early endocytic pathway. — *J. Cell Sci.* **117**: 6297–6311.
- STENMARK H. 2009: Rab GTPases as coordinators of vesicle traffic. — *Nat. Rev. Mol. Cell Biol.* **10**: 513–525.
- ULLRICH O., REINSCH S., URBE S., ZERIAL M. & PARTON R.G. 1996: Rab11 regulates recycling through the pericentriolar recycling endosome. — *J. Cell Biol.* **135**: 913–924.
- UNO T., HATA K., HIRAGAKI S., ISOYAMA Y., TRANGLE T.D., UNO Y., KANAMARU K., YAMAGATA H., NAKAMURA M. & TAKAGI M. 2010: Small GTPases of the Rab family in the brain of *Bombyx mori*. — *Histochem. Cell Biol.* **134**: 615–622.
- UNO T., SAKAMOTO K., ISOYAMA Y., HIRAGAKI S., UNO Y., KANAMARU K., YAMAGATA H., TAKAGI M., MIZOGUCHI A. & TAKEDA M. 2013: Relationship between the expression of Rab family GTPases and neuropeptide hormones in the brain of *Bombyx mori*. — *Histochem. Cell Biol.* **139**: 299–308.
- UNO T., ISOYAMA Y., SAKAMOTO K., UNO Y., SAKAMOTO K., KANAMARU K., YAMAGATA H., TAKAGI M., MIZOGUCHI A. & TAKEDA M. 2014: Characterization of Rab-interacting lysosomal protein in the brain of *Bombyx mori*. — *Histochem. Cell Biol.* **141**: 311–320.
- UNO T., FURUTANI M., WATANABE C., SAKAMOTO K., UNO Y., KANAMARU K., YAMAGATA H., MIZOGUCHI A. & TAKEDA M. 2016: Rab proteins in the brain and corpus allatum of *Bombyx mori*. — *Histochem. Cell Biol.* **146**: 59–69.
- UNO T., FURUTANI M., SAKAMOTO K., UNO Y., KANAMARU K., MIZOGUCHI A., HIRAGAKI S. & TAKEDA M. 2017: Localization and functional analysis of the insect-specific RabX4 in the brain of *Bombyx mori*. — *Arch. Insect Biochem. Physiol.* **96**: e21404, 13 pp.
- UNO T., OZAKIYA Y., FURUTANI M., SAKAMOTO K., UNO Y., KAJIWARA H., KANAMARU K. & MIZOGUCHI A. 2019: Functional characterization of insect-specific RabX6 of *Bombyx mori*. — *Histochem. Cell Biol.* **151**: 187–198.
- UNO T., OZAKIYA Y., SASAO M., SAKAMOTO K., YAMAUCHI Y., UNO Y., KANAMARU K. & MIZOGUCHI A. 2021: Relationship between Rab and insulin-like proteins in the nervous system of *Bombyx mori*. — *Eur. J. Entomol.* **118**: 307–314.
- UYTTERHOEVEN V., KUENEN S., KASPROWICZ J., MISKIEWICZ K. & VERSTREKEN P. 2011: Loss of skywalker reveals synaptic endosomes as sorting stations for synaptic vesicle proteins. — *Cell* **145**: 117–132.
- VAFOPOULOU X. & STEEL C.G. 2014: Synergistic induction of the clock protein PERIOD by insulin-like peptide and prothoracicotrophic hormone in *Rhodnius prolixus* (Hemiptera): implications for convergence of hormone signaling pathways. — *Front. Physiol.* **5**: 41, 12 pp.
- VAFOPOULOU X., STEEL C.G. & TERRY K.L. 2007: Neuroanatomical relations of prothoracicotrophic hormone neurons with the circadian timekeeping system in the brain of larval and adult *Rhodnius prolixus* (Hemiptera). — *J. Comp. Neurol.* **503**: 511–524.
- VILLALOBOS-SAMBUCARO M.J., LORENZO-FIGUEIRAS A.N., RICCILO F.L., DIAMBRA L.A., NORIEGA F.G. & RONDEROS J.R. 2015: Allatotropin modulates myostimulatory and cardioacceleratory activities in *Rhodnius prolixus* (Stal). — *PLoS ONE* **10**(4): e0124131, 14 pp.
- WANG T., MING Z., XIAOCHUN W. & HONG W. 2011: Rab7: role of its protein interaction cascades in endo-lysosomal traffic. — *Cell Signal* **23**: 516–521.
- WANG Q., MOHAMED A.A. & TAKEDA M. 2013: Serotonin receptor B may lock the gate of PTTH release/synthesis in the Chinese silkworm, *Antheraea pernyi*: a diapause initiation/maintenance mechanism? — *PLoS ONE* **8**(11): e79381, 13 pp.
- WISE S., DAVIS N.T., TYNDALE E., NOVERAL J., FOLWELL M.G., BEDIAN V., EMERY I.F. & SIWICKI K.K. 2002: Neuroanatomical studies of period gene expression in the hawkmoth, *Manduca sexta*. — *J. Comp. Neurol.* **447**: 366–380.

- XU G., TENG Z.W., GU G.X., QI Y.X., GUO L., XIAO S., WANG F., FANG Q., WANG F., SONG Q.S. ET AL. 2020: Genome-wide characterization and transcriptomic analyses of neuropeptides and their receptors in an endoparasitoid wasp, *Pteromalus puparum*. — *Arch. Insect Biochem. Physiol.* **103**(2): e21625, 23 pp.
- YE T., TANG W. & ZHANG X. 2012: Involvement of Rab6 in the regulation of phagocytosis against virus infection in invertebrates. — *J. Proteome Res.* **11**: 4834–4846.
- ZENG H., QIN Y., DU E., WEI Q., LI Y., HUANG D., WANG G., VEENSTRA J.A., LI S. & LI N. 2021: Genomics- and peptidomics-based discovery of conserved and novel neuropeptides in the American cockroach. — *J. Proteome Res.* **20**: 1217–1228.

Received November 23, 2022; revised and accepted March 14, 2023
Published online March 29, 2023