

The effects of Pleistocene glaciations on the phylogeography of *Melitaea cinxia* (Lepidoptera: Nymphalidae)

NIKLAS WAHLBERG^{1*} and ILIK SACCHERI²

¹Department of Zoology, Stockholm University, 10691 Stockholm, Sweden

²School of Biological Sciences, University of Liverpool, The Biosciences Building, Crown Street, Liverpool L69 7ZB, UK

Key words. Phylogeography, Palearctic, cytochrome oxidase I, Lepidoptera

Abstract. Partial (600 bp) sequences of mitochondrial cytochrome oxidase I (COI) gene were used to infer the phylogeography of *Melitaea cinxia* (Lepidoptera: Nymphalidae) across the entire distributional range of the species, encompassing north Africa and Eurasia. Cladistic analysis of 49 distinct haplotypes (haplotype and nucleotide diversity were 0.95 and 0.027, respectively) revealed strong phylogeographic structure in *M. cinxia*, characterised by four major clades: Morocco; Western (Iberia, France, Italy); Central (central and northern Western Europe, Balkans, Greece, Anatolia, Levant); and Eastern (eastern Baltic, Urals, Iran, Siberia, China); separated by average pairwise distances of between 2 and 6 percent. This pattern is consistent with the location of southern glacial refugia in the Iberian, Italian and Balkan peninsulas, as well as multiple eastern refugia. The Western clade is further structured into south-central Iberian, northern Iberian (and French) and southern Italian sub-clades; and the Eastern clade into Near Eastern and Far Eastern sub-clades; with weaker phylogeographical concordance within the Central clade, except for a large area in central and northern Western Europe which is monomorphic for COI haplotype. The Baltic and eastern Europe have been primarily colonized by the Far Eastern sub-clade, rather than the Central (Balkan) clade, highlighting the importance of including Near and Far Eastern populations in phylogeographic studies of Palearctic species. Maps showing the extent of clades and sub-clades suggest several regions of secondary contact and possible hybridization. Interspecific comparison of representative *M. cinxia* haplotypes supports a monophyletic origin of all *M. cinxia*.

INTRODUCTION

Quaternary climatic oscillations, punctuated by the Pleistocene glaciations, caused massive changes to the distributions of species in the Palearctic (Hewitt, 2000, 2004; Schmitt, 2007). Repeated cycles of demographic contraction and expansion, into and out of pockets of climatically protected regional refugia, combined with individual ecologies, has led to varied patterns of present day phylogeographic concordance among species (Hewitt, 1999). These patterns indicate the existence of multiple glacial refugia, large differences in the ages of intraspecific divergences, and differential rates of post-glacial expansion among refugial populations. Isolation among refugial populations promotes genetic and phenotypic differentiation as a result of independent adaptation to local environments and genetic drift, with consequences for reproductive isolation between discrete refugial lineages and the creation of hybrid zones where diverged lineages come into secondary contact (Hewitt, 1999). Thus, the Pleistocene glaciations are likely to have increased the effective size of many populations at the species level, but the same process typically reduces genetic diversity at the regional level, usually in proportion to the distance from the parent refugium (e.g. Bernatchez & Wilson, 1998; Schmitt & Seitz, 2002; Michaux et al., 2003) and depending on the mode of contraction and expansion (Ibrahim et al., 1996).

Phylogeographic studies of Palearctic species are burgeoning but it remains the case that current phylogeographic paradigms for Europe are limited by the relatively small number of comprehensive case studies (Hewitt, 1999, 2004). Moreover, the overwhelming majority of intraspecific phylogeographic studies of Palearctic species have focused on European populations of species that often have much wider distributions (Taberlet et al., 1998; Hewitt, 1999), meaning that our understanding of the history of Palearctic species is based upon patterns found in only one-fifth of the landmass of this biogeographical region. This is especially a problem for insects, for which the fossil and subfossil record is very poor (with the exception of beetles), and for which inference of evolutionary history is currently based entirely on molecular data, particularly DNA sequences from the mitochondrial genome. Butterflies are no exception and the few published studies have been entirely restricted to western Europe (e.g., Joyce & Pullin, 2001; Schmitt & Seitz, 2001; Aagaard et al., 2002; Schmitt & Seitz, 2002; Habel et al., 2005; Schmitt et al., 2005; Weingartner et al., 2006), with the exception of one study on *Aglais urticae* (Vandewoestijne et al., 2004), and another on *Hesperia comma* (Forister et al., 2004) focused on North America but with some sampling of eastern and western Eurasia.

* Corresponding and current address: Laboratory of Genetics, Department of Biology, University of Turku, 20014 Turku, Finland; e-mail: niklas.wahlberg@utu.fi

We take a phylogeographic approach (Avise, 2000) to investigate the biogeographic history of the ecologically well-studied butterfly *Melitaea cinxia* (L.), the Glanville fritillary. *M. cinxia* is found on dry meadows containing its host plants *Plantago* and *Veronica* (both in Plantaginaceae) from the Atlantic coast to the Baikal Lake region in Siberia (Hanski, 1999) and to northwestern China (I. Hanski, pers. comm.). It is distributed between approximately 35°N and 60°N, though in the southern parts of its range it is restricted to higher elevations. There is a population in the Atlas Mountains of Morocco in northern Africa, which is often considered to be subspecifically distinct (Tolman & Lewington, 1997). Otherwise, *M. cinxia* is known for its morphological invariability. Although some populations have been described as subspecifically distinct (Tuzov, 2000; Churkin & Koleznichenko, 2002), many authors consider all Eurasian populations of *M. cinxia* to belong to the nominotypical subspecies (e.g. Higgins, 1941; Tolman & Lewington, 1997).

MATERIAL AND METHODS

We sampled *Melitaea cinxia* from as wide a range of localities as possible (Table 1). Individuals were mainly sampled as adults, but in some cases larvae were collected and either reared to adults or used directly in DNA extraction. We extracted DNA mainly from two legs of dried specimens according to protocols described previously (Zimmermann et al., 2000; Wahlberg et al., 2003). We amplified the first half (600 bp) of the cytochrome oxidase subunit I (COI) gene in the mitochondrial genome from all the sampled individuals using the primer pair LCO and HCO (Folmer et al., 1994). The cycling profile for the PCR reactions was 95°C for 2 min, 35 cycles of 94°C for 30 s, 47°C for 30 s, 72°C for 1 min and a final extension period of 72°C for 10 min. PCR fragments were then sequenced using either an ABI 377 Automated Sequencer or a Beckman-Coulter CEQ8000 capillary sequencer (Bromma, Sweden) and a dye terminator cycle sequencing kit. Sequences were aligned by eye as the length of this protein-coding gene is highly conserved and insertions or deletions have not been observed in melitaeines (Wahlberg & Zimmermann, 2000; Zimmermann et al., 2000). Each unique haplotype sequence is available from GenBank (Table 1).

Initially all the *M. cinxia* sequences were analyzed together to identify unique haplotypes in the program Arlequin 3 (Excoffier et al., 2005). The phylogenetic relationships of the unique haplotypes were then analyzed using parsimony and rooting with the outgroup *Melitaea aurelia* (Wahlberg & Zimmermann, 2000; Wahlberg et al., 2005b). Most parsimonious cladograms were searched for from the equally weighted and unordered data matrix using a heuristic search algorithm of NONA 2.0 (Goloboff, 1998). The heuristic searches were conducted with 10-1000 random addition replicates using tree bisection-reconnection (TBR) branch swapping with up to twenty trees held during each step. Trees were drawn with the aid of Winclada (Nixon, 2002). Clade support was evaluated using bootstrap with 1000 pseudoreplicates, with 5 random addition replicates within each pseudoreplicate. In addition, a minimum-spanning network was constructed for all individuals using the program Arlequin 3 (Excoffier et al., 2005).

The patterns of nucleotide substitutions among sequences were analyzed using the program Arlequin 3 (Excoffier et al., 2005) and MEGA3 (Kumar et al., 2004). Standard genetic

indices [haplotype diversity (h) and nucleotide diversity (π)] were computed.

To verify the monophyly of *M. cinxia* haplotypes, a subset of ten haplotypes, representing the main lineages, was analysed with all the homologous *Melitaea* sequences from the study by Wahlberg & Zimmermann (2000). The resulting *Melitaea* tree was rooted using the related *Chlosyne acastus* from the same study.

RESULTS

The 120 individuals of *M. cinxia* from 58 localities yielded 49 haplotypes, of which 34 were found only in one individual (Table 1). The 600 bp showed 77 polymorphic sites with 71 observed transitions and 11 observed transversions. Haplotype diversity (h) was 0.95 (± 0.01) and nucleotide diversity (π) was 0.027 (± 0.013). The uncorrected pairwise differences of the haplotypes varied widely from 0.1% to more than 6%, with the three haplotypes from Morocco being most diverged from the others (Table 2). One haplotype (Haplotype 25, Table 1) was particularly widespread, being found in Ukraine, northern Italy, Denmark, Sweden, Poland, Belgium and northern France. A few other haplotypes were found in more than one collection locality, although generally from localities relatively close to each other. Haplotype 4 was found in southern, central and northern Spain; haplotype 19 in northern Spain and eastern France; and haplotype 43 in Saratov and northern Iran. The greatest geographical separation between two identical haplotypes was for haplotype 31, found in the Lake Baikal region and in Åland, Finland.

Cladistic analysis of the *Melitaea* sequences yielded 28 equally parsimonious trees (length 499 steps, CI = 0.48, RI = 0.66), of which the strict consensus is shown (Fig. 1). The subset of ten *M. cinxia* haplotypes formed a strongly supported monophyletic group (bootstrap value = 100%). Within *M. cinxia*, the representative Moroccan haplotype (haplotype 1) was clearly sister to the rest of the haplotypes (bootstrap value = 72%).

Of the 77 variable sites, 44 were parsimony informative within *M. cinxia*. Cladistic analysis of the 49 *M. cinxia* haplotypes resulted in 408 equally parsimonious trees (length 178 steps, CI = 0.68, RI = 0.89). The low level of homoplasy (as shown by the CI and RI values) indicates that there is strong structure in the data set. Indeed the resulting trees are impervious to assumptions of transition:transversion weightings. Of the twelve inferred transversions ten are synapomorphic (or autapomorphic) with no reversals. Two are homoplastic and occur independently in different clades. Weighting transversions up to 10 times transitions has no effect on the topology or number of trees found.

There are four major clades in the haplotype tree (Fig. 2), which correspond to clearly definable geographical areas (Fig. 3). The Western clade contains haplotypes found mainly on the Iberian peninsula but also in France, Italy and Isle of Wight. The Central clade is distributed throughout central Europe from southern Sweden down to northern Italy and across to Turkey and Lebanon. The Eastern clade stretches out from central Sweden across

TABLE 1. Geographical origin and collection year of *Melitaea cinxia* individuals used in this study, listed in order of COI haplotype number.

Haplotype i.d.	Collection locality	Year	Voucher code	GenBank Accession
1	MOROCCO: Haut Atlas septentrional, Cirque de Jaffar, 2000–2880 m, Djebel Ayachi, SW Midelt	2002	NW103-9	EF680408
1	MOROCCO: Moyen Atlas central, Col du Zad, 1800–2500 m, S Timahdite	2002	NW103-10	
1	MOROCCO: Moyen Atlas central, Tizi-n-Tretten, 1900–2000 m, SE Ifrane	2002	JL3-1	
2	MOROCCO: Moyen Atlas central, Tizi-n-Tretten, SE Ifrane	2002	NW103-11	EF680409
3	MOROCCO: Moyen Atlas central, Tizi-n-Tretten, 1900–2000 m, SE Ifrane	2002	JL3-2	EF680410
4	SPAIN: Catalonia	–	AC6-5	
4	SPAIN: Catalonia	–	AC7-1	
4	SPAIN: Catalonia, Vallès Oriental, El Puig	1998	NW34-14	EF680411
4	SPAIN: P.N. Cazorla Segura, Arroyo Frio	1997	IS S2	
4	SPAIN: P.N. Cazorla Segura, Arroyo de Las Truchas	1995	IS S3	
4	SPAIN: P.N. Cazorla Segura, Arroyo Frio	1997	IS S5	
4	SPAIN: Navalguijo (Avila), Sierra de Gredos	1986	SV1	
4	SPAIN: Valsain, Sierra de Guadarrama (Segovia)	1993	SV2	
4	SPAIN: Sierra de Alcaraz (Albacete)	1989	SV4	
4	SPAIN: Sierra de Alcaraz (Albacete)	1988	SV5	
4	SPAIN: Sierra de Alcaraz (Albacete)	1989	SV6	
4	SPAIN: Navalguijo (Avila), Sierra de Gredos	1986	SV7	
5	FRANCE: Mourmelon, N de Châlons	1998	IS F5	EF680412
6	SPAIN: Madrid, Miraflores	1999	S6	EF680413
6	SPAIN: Manzanares el Real, Madrid	1990	SV3	
6	SPAIN: Madrid	1998	SV8	
7	ITALY: Sicilia, Madonie	1998	NW28-6	EF680414
8	FRANCE: Cervieres, SE of Briançon	1998	C5	EF680415
9	ITALY: Emilia Romagna, Piacenza, Piani di Aglio (900m)	1995	I-1	EF680416
9	ITALY: Emilia Romagna, Piacenza, Piani di Aglio (900m)	1995	I-3	
10	ITALY: Calabria, Catanzaro	1998	NW28-8	EF680417
11	ITALY: Lazio, Riofreddo	1998	NW28-9	EF680418
12	ITALY: Abruzzo, Capistrello	1998	NW28-10	EF680419
13	FRANCE: Montpellier		NW F9-2	EF680420
14	ITALY: Liguria, Imperia, Carmo Langan (1120)	1994	I-4	EF680421
14	ITALY: Liguria, Imperia, Carmo Langan (1120)	1994	I-5	
15	FRANCE: Montpellier	1997	NW F9-1	EF680422
16	FRANCE: Massignieu	1997	NW F7-4	EF680423
17	FRANCE: Mourmelon, N de Châlons	1997	NW35-2	EF680424
18	SPAIN: Madrid, Montejo de la Sierra	1983	NW35-3	EF680425
19	SPAIN: Alt Empordà, Catalonia, Darnius	1998	NW34-15	EF680426
19	FRANCE: Massignieu	1997	NW F7-1	
19	FRANCE: Massignieu	1997	NW F7-3	
20	UK: Isle of Wight, Compton	1999	UK1_1_1	EF680427
20	UK: Isle of Wight, Chilton	1999	UK2_1_1	
20	UK: Isle of Wight, Chilton	1999	UK2_2_1	
20	UK: Isle of Wight, Chilton	1999	UK2_3_1	
20	UK: Isle of Wight, Chilton	1999	UK2_4_1	
20	UK: Isle of Wight, Sheppards	1999	UK3_1_1	
20	UK: Isle of Wight, Sheppards	1999	UK3_2_1	
20	UK: Isle of Wight, Sheppards	1999	UK3_3_1	
20	UK: Isle of Wight, Old Park	1999	UK4_1_1	
20	UK: Isle of Wight, Old Park	1999	UK4_2_1	
20	UK: Isle of Wight, Old Park	1999	UK4_3_1	
21	LEBANON: Mohafazat Zahlé, Bawarij	1998	NW34-13	EF680428
22	ALBANIA: Mt. Dajti	2006	NW159-1	EF680429
23	TURKEY: Sultan Daglari, Aksehir	1997	NW19-4	EF680430
24	BULGARIA: Rhodopi, Lukovitsa Lives	1993	NW30-3	EF680431
25	GERMANY: Thüringen, (45km NW of Bamberg)	1999	SV12	
25	UKRAINE	–	NW22-6	EF680432
25	ITALY: Florence	1998	NW28-7	
25	POLAND: Krakow, Klucze	1999	P12	

TABLE 1 continued.

Haplotype i.d.	Collection locality	Year	Voucher code	GenBank Accession
25	POLAND: Krakow, Klucze	1999	P13	
25	BELGIUM: Military Camp of Lagland (SW of Arlon)	1996	B1	
26	GREECE: Peloponnesos, Diakofto	1998	NW30-5	EF680433
27	GREECE: Pissoderi	1997	NW23-13	AF187753
28	HUNGARY: Jósvalfö	1996	NW28-4	EF680434
29	TURKEY: Ürgüp	1997	NW19-3	EF680435
30	RUSSIA: Buryatia, Utitzcina	1998	NW26-7	EF680436
31	RUSSIA: Buryatia, Ubukun	1998	NW25-2	EF680437
31	FINLAND: Åland	1995	Fin200	
32	FINLAND: Houtskär	1998	NW35-1	EF680438
32	FINLAND: Åland	1995	Fin168	
32	FINLAND: Åland	1995	Fin239	
32	FINLAND: Åland	1995	Fin276	
32	FINLAND: Åland	1995	Fin713	
32	FINLAND: Åland	1995	Fin812	
32	FINLAND: Åland	1995	Fin901	
32	FINLAND: Åland	1995	Fin966	
32	FINLAND: Åland	1995	Fin1154	
32	FINLAND: Åland	1995	Fin1185	
32	FINLAND: Åland	1995	Fin1233	
32	FINLAND: Åland	1995	Fin1287	
32	SWEDEN: Väddö, Edenby	2006	SW1	
32	SWEDEN: Väddö, Harsängen	2006	SW4	
33	FINLAND: Åland	1995	Fin747	EF680439
33	FINLAND: Åland	1995	Fin1563	
34	LITHUANIA: Kaulakiai, Raseiniai	2001	SV18	EF680440
35	ESTONIA: Paldiski	1998	NW33-4	EF680441
36	FINLAND: Åland	1995	Fin22	EF680442
36	FINLAND: Åland	1995	Fin1075	
36	FINLAND: Åland	1995	Fin1397	
36	FINLAND: Åland	1995	Fin1545	
36	SWEDEN: Väddö, Norrbylle Mellesta	2006	SW2	
36	SWEDEN: Väddö, Vorrund	2006	SW3	
37	SWEDEN: Stockholm archipelago	2001	NW73-14	AY788656
38	LITHUANIA: Kaulakiai, Raseiniai	2000	SV17	EF680443
38	POLAND: Krakow, Klucze	1999	P2	
38	POLAND: Krakow, Klucze	1999	P5	
38	POLAND: Krakow, Klucze	1999	P7	
38	POLAND: Krakow, Klucze	1999	P8	
38	POLAND: Krakow, Klucze	1999	P9	
38	POLAND: Krakow, Klucze	1999	P10	
38	POLAND: Krakow, Klucze	1999	P11	
38	POLAND: Krakow, Klucze	1999	P14	
39	CHINA: Xinjiang, Tian Shan	2002	NW96-1	EF680444
40	RUSSIA: Buryatia, Djirga	1998	NW28-11	EF680445
41	ESTONIA: Hiiumaa	1994	NW ee	EF680446
42	RUSSIA: Ural Mountains	1999	SV15	EF680447
43	RUSSIA: Saratov, Saratov	1997	NW15-15	EF680448
43	IRAN: Ardabil	2003	NW120-13	
44	IRAN: Tehran, Elburs-Gebirge	2003	NW120-12	EF680449
45	RUSSIA: Ural Mountains	1999	SV13	EF680450
46	LITHUANIA: Kaulakiai, Raseiniai	1999	SV16	EF680451
46	POLAND: Krakow, Klucze	1999	P1	
46	POLAND: Krakow, Klucze	1999	P4	
46	SWEDEN: Grasö, Norrboda	2006	SW5	
47	POLAND: Krakow, Klucze	1999	P6	EF680452
48	RUSSIA: Ural Mountains	1999	SV14	EF680453
49	RUSSIA: S Ural Mountains	1999	AC7-16	EF680454

TABLE 2. Average (\pm SE) uncorrected pairwise distances of cytochrome oxidase subunit I (COI) haplotypes for the major *Melitaea cinxia* lineages.

	Morocco	West	Central	East
Morocco	0.004 \pm 0.002	0.050 \pm 0.008	0.052 \pm 0.008	0.062 \pm 0.009
West		0.013 \pm 0.003	0.023 \pm 0.005	0.040 \pm 0.007
Central			0.007 \pm 0.002	0.032 \pm 0.006
East				0.011 \pm 0.002

Finland to the Baikal Lake region and down into north-western China and northern Iran. The haplotypes found in Morocco form their own lineage.

The Eastern clade appears to be the sister group of the Central clade, with the Western clade being sister to these two in the most parsimonious reconstruction of the data set. While this arrangement receives only moderate (79%) bootstrap support, the sister relationship of the Eastern and Central clades is supported by a unique nonsynonymous change at site 505 which leads to an amino acid change of an isoleucine to a valine. There are 8 other inferred amino acid changes, but these are autapomorphic or define small clades of two or three haplotypes. It can be noted that a neighbour joining algorithm using distances estimated with a variety of models (uncorrected *p*-values, the HKY85 model and the LogDet model) places the Western and Central clades as sister groups. However, such a scenario would entail independent acquisition of the amino acid change described above, or alternatively a single acquisition and a subsequent reversal to isoleucine in the Western clade. We prefer the more parsimonious scenario of a single acquisition of the amino acid change with no subsequent reversal.

A minimum-spanning network of the *M. cinxia* haplotypes (Fig. 4) shows the same patterns as the phylogeny, with the four major groups of haplotypes (Morocco, Western, Central and Eastern) clearly separated from each other by 9 or more inferred changes. Because of unequal sampling effort among localities the size of haplotype nodes in Fig. 4 does not accurately reflect the abundance of each haplotype in *M. cinxia* at the species level. This caveat applies principally to haplotypes 20, 32 and 38, respectively from Isle of Wight, Åland and Klucze, the three largest single locality samples. On the other hand, the large nodes for haplotypes 4 and 25, which include samples from many geographically separated localities, are genuinely common haplotypes at the species level.

The minimum-spanning network (Fig. 4) and sub-clades in Fig. 2 reveal further phylogeographic structuring within the major clades, particularly the Western and Eastern clades. A colour-coded map (Fig. 5) shows the distribution of these sub-clades. The Western clade has three distinct groups which we designate “Iberia”, “France” and “southern Italy”, with France being the sister group to Iberia (Fig. 2). Samples in the Iberia sub-clade originate almost exclusively from Spain, whereas the France sub-clade is predominantly composed of samples from southern and eastern France but also includes samples from the northeastern corner of Spain (and one central Spanish sample), the coastal Italy-France border,

and the Isle of Wight. The southern half of Italy, including Sicily, appears to be monophyletic at the sub-clade level, though two samples falling into the southern Italian sub-clade originate from northern Italy and the northeast Italy-France border.

The Eastern clade consists of two geographically structured sub-clades, one occurring predominantly in the southwestern distribution of the clade (“Ural” sub-clade), and the other covering a very extensive area from the far east to the northwest of the clade (“Siberia” sub-clade). The Ural sub-clade contains haplotypes from the southern Ural Mountains, Saratov (Russia), northern Iran, Poland, Lithuania and the island of Hiiumaa (Estonia); the Siberia sub-clade contains haplotypes from Buryatia (Lake Baikal region), northwestern China and, surprisingly, the northern Estonian coast, Åland (Finland) and the Stockholm archipelago (Sweden). The Central clade does not exhibit any strong internal structure (Fig. 4) and shows the least amount of variation (Table 2). Fig 3 indicates regions of secondary contact between the major clades;




Fig. 1. Strict consensus of 28 most parsimonious trees (length 539 steps, CI = 0.47, RI = 0.81) inferred for 600 bp of COI gene. Numbers above branches are bootstrap proportions.




Fig. 2. Strict consensus of 408 most parsimonious trees for *Melitaea cinxia* haplotypes inferred for 600 bp of COI gene. Major lineages coded with symbols: square = Morocco, triangle = Western clade, star = Central clade, and circle = Eastern clade. Numbers above branches are bootstrap proportions.

Fig. 5 additionally gives an impression of where secondary contact between sub-clades is likely to occur.

DISCUSSION

The COI haplotypes sequenced for this study support a monophyletic origin of all *Melitaea cinxia*, and have revealed a high degree of phylogeographic structuring across the species range. Taking the evidence given in Fig. 2, the deepest and earliest divergence occurred between Eurasian populations and populations in the

North African Atlas mountains. The next split is likely to have occurred between the ancestors of the Western clade and Central + Eastern clades, closely followed by the final major divergence between the ancestors of Central and Eastern clade populations. Interestingly, the deep split between European and North African populations has been found in other butterfly species, such as *Pararge aegeria* (Weingartner et al., 2006) and two species of *Polyommatus* (Wiemers & Fiedler, 2007). However, the COI distance between the two populations of *P. aegeria*




Fig. 3. Map showing localities of origin of specimens. Symbol codes according to Fig. 2: square = Morocco, triangle = Western clade, star = Central clade, and circle = Eastern clade. Overlapping symbols indicate where more than one major clade is represented in the same locality.




Fig. 4. Minimum Spanning network of *Melitaea cinxia* haplotypes. Size of circles proportional to number of individuals with that haplotype. Numbers within or next to circles are the haplotype number. Black dots along lines indicate number of nucleotide substitutions separating connected haplotypes. Colour codes indicate COI haplotype clades or sub-clades: black = Morocco clade (H1–H3), dark blue = Iberia sub-clade (H4–H6), light blue = France sub-clade (H13–H20), purple = southern Italy sub-clade (H7–H12), green = Central clade (H21–H29), orange = Urals sub-clade (H41–H49), yellow = Siberia sub-clade (H30–H40).




Fig. 5. Geographic distribution of COI sub-clades in Europe, as indicated by haplotype tree (Fig. 2) and minimum-spanning network (Fig. 4). Colour codes are the same as in Fig. 4: dark blue = Iberia sub-clade, light blue = France sub-clade, purple = southern Italy sub-clade, green = Central clade, orange = Urals sub-clade, yellow = Siberia sub-clade. Multi-coloured circles indicate the presence of haplotypes from more than one clade or sub-clade at the same locality. The sub-clade designation of eastern samples not included in this map is Urals for Urals and Iran, and Siberia for Baikal and China.

is less than half (about 2%, Weingartner et al., 2006) the distance we have found for *M. cinxia* (more than 5%, Table 2).

Weingartner et al. (2006) suggest that the ancestor of the genus *Pararge* colonised North Africa over 5 million years ago and subsequently recolonised Europe about 1 million years ago. In the case of *M. cinxia*, we cannot say much about the direction of colonisation since it may be that there is no extant sister species of *M. cinxia*, but rather that *M. cinxia* is sister to a large clade of Eurasian species related to *M. diamina*, *M. arcesia* and *M. athalia* (Wahlberg & Zimmermann, 2000; Wahlberg et al., 2005b). In any case, it appears that the split between populations in North Africa and Eurasia occurred about twice as much earlier in *M. cinxia* than in *P. aegeria* (assuming a similar rate of molecular evolution in COI). Alternatively, the rate of molecular evolution could be very different in these two species of Nymphalidae, leading to different estimates of relative ages of divergence. Absolute times of divergence in *M. cinxia* can only be investigated in a study of the entire genus *Melitaea* using age estimates from a recent study on the subfamily Nymphalinae (Wahlberg, 2006).

The Western and Central clades correspond well to having originated in the previously hypothesized glacial refugia in the Iberian, Italian, and Balkan Peninsulæ (Hewitt, 1999). In the Western clade, all the populations

from southern Italy appear to be sister to the populations from the Iberian peninsula and France, but in northern Italy haplotypes from France sub-clade, Central clade, and southern Italy sub-clade are all present. This pattern of haplotypes in Italy suggests that the characteristic southern Italian sub-clade diverged from the Iberian-French lineage in a southern Italian refugium during one or more glacials, and has subsequently expanded northwards, while French and Balkan haplotypes have colonized northern Italy from the west and east respectively. The other division of the Western clade into Iberia and France sub-clades supports the existence of two separate refugia for *M. cinxia* in southwestern Europe, one in southern Spain and another along coastal regions of southern France or northeastern Spain, consistent with mounting evidence of multiple glacial refugia in Iberia (Bella et al., 2006; Gómez & Lunt, 2007; Schmitt, 2007).

The geographic position of the Central clade is concordant with the idea of a Balkan refuge, although additional more southerly refuges in Greece, Turkey, and the Levant cannot be ruled out. Although there is much less divergence among COI lineages in the Central clade compared to the other two major clades, there is a degree of phylogenetic separation between haplotype 25 from central and northwestern Europe (northern France, Germany, Belgium, northern Italy, Denmark, southern Sweden, Poland, Hungary, Ukraine) and the remaining Central clade hap-

lotypes in southeastern Europe, indicating that central and northwestern Europe was colonized by a single, probably Balkan lineage, while southeastern populations are derived from a diversity of refugial lineages.

The role of temperate Asia in the Pleistocene biogeographical history of Palaearctic insect species has not been studied in detail. Our results indicate that the area has been important in the evolutionary history of *M. cinxia*. Our sampling is rather sparse compared to European populations (Fig. 3), yet we find internal phylogeographic structure in the Eastern clade (Fig. 2), corresponding to a Caspian Sea refuge and a Far Eastern refuge. Given the relative geographical proximity of the Baltic region to the European refugia, it was a surprise to find that current populations of *M. cinxia* in Åland, Finland, and eastern Sweden are closely related to Far Eastern populations, in contrast to several other animal and plant species in this region that are typically derived from a Balkan refugium (Hewitt, 1999). Possible explanations include a faster rate of post-glacial expansion from a Far Eastern refugium, associated with a more continuously favourable habitat, perhaps combined with pre-adaptation to colder northern climates. Another interesting aspect of the Far Eastern expansion is the retention of genetic diversity at what we presume to be the northwestern colonizing front of this clade. The sparse sampling across Near and Far Eastern Asia does not allow critical assessment of haplotype diversity along hypothetical colonization routes, but the presence of eight distinct Siberia clade haplotypes in the Baltic region implies a relatively even demographic expansion. This pattern of mitotype diversity is in marked contrast to the significantly reduced COI variation in north European populations of the saproxylic beetle, *Phyto kolwensis* (Painter et al., 2007), also derived from eastern Eurasia. The far greater extent, during the last glacial maximum, of steppe habitat suitable for *M. cinxia*, compared to the much more restricted distribution of old spruce forest, on which *P. kolwensis* is dependent, is likely to have been a key factor underlying this difference (Grichuk, 1984; Tarasov et al., 2000).

The haplotype maps (Fig. 3 and 5) give some idea of the position of secondary contact zones between the major clades, and also between sub-clades. The Western (both the Iberia and France sub-clades) and Central clades meet in northeastern France and northern Italy (southern Italy and France sub-clades), where contact between France and southern Italy sub-clades is also likely. The main regions of secondary contact between Iberia and France sub-clades are in Catalonia and central Spain. In eastern Europe we detected Central and Eastern (Siberia and Urals sub-clade) haplotypes in one Polish population, and haplotypes from both Eastern sub-clades in Lithuania, Estonia, and eastern Sweden. Eastern Ukraine, eastern Turkey, and Kazakhstan, where we have not sampled, represent further regions of potential secondary contact. This study of mitochondrial haplotypes gives no information on the extent of hybridization and nuclear introgression between the clades and sub-clades, but pro-

vides a basis for designing hybrid zone studies, including fitness consequences of hybridization in this species.

M. cinxia is a widely distributed species but has suffered severe declines in western Europe since the 1960s, largely as a result of agricultural intensification and urbanization (Hanski & Kuussaari, 1995). In this context, this phylogeographic study has some bearing on the conservation of this species, particularly as regards phylogenetic uniqueness of regional populations and reintroduction or augmentation programmes. For example, we note that the remnant UK population on the Isle of Wight is monomorphic for COI and is derived from the France sub-clade, but that surviving Belgian and German populations belong to the Central clade. These observations suggest that the now extinct populations in mainland UK and Holland were derived from France sub-clade and Central clade, respectively.

The COI haplotype variation in *M. cinxia* surpasses any other Eurasian butterfly species studied to date (Vandewoestijne et al., 2004; Wahlberg et al., 2005a; Weingartner et al., 2006). In contrast to the highly mobile *Aglaia urticae* (Vandewoestijne et al., 2004), this variation is shown to have a strong phylogeographic signal, consistent with short-range dispersal (Hanski et al., 1994; Hanski, 1999) and host plant specificity (Kuussaari et al., 2000) in *M. cinxia*. Our results suggest that *M. cinxia* has had continuously large, widespread populations over the past 2 million years, and that local population dynamics (Hanski, 1999; Ehrlich & Hanski, 2004), coupled with multiple isolated glacial refugia, have allowed many haplotypes to persist through repeated cycles of demographic expansion and contraction.

ACKNOWLEDGEMENTS. A significant impediment to realizing large-scale phylogenies is the collection of samples from across species ranges. We are indebted to the many people, too numerous to list here, who contributed samples for this study. We thank I. Hanski for support and patience throughout this longterm project, T. Schmitt and an anonymous reviewer for useful suggestions on an earlier draft, and J. Leneveu for help in the lab.

REFERENCES

- AAGAARD K., HINDAR K., PULLIN A.S., JAMES C.H., HAMMARSTEDT O., BALSTAD T. & HANSEN O. 2002: Phylogenetic relationships in brown argus butterflies (Lepidoptera: Lycaenidae: Aricia) from north-western Europe. *Biol. J. Linn. Soc.* **75**: 27–37.
- AVISE J.C. 2000: *Phylogeography: The History and Formation of Species*. Harvard University Press, Cambridge, MA, 447 pp.
- BELLA J.L., SERRANO L., ORELLANA J. & MASON P.L. 2006: The origin of the *Chortippus parallelus* hybrid zone: chromosomal evidence of multiple refugia for Iberian populations. *J. Evol. Biol.* **20**: 568–576.
- BERNATCHEZ L. & WILSON C.C. 1998: Comparative phylogeography of nearctic and palaearctic fishes. *Mol. Ecol.* **7**: 431–452.
- CHURKIN S.V. & KOLESNICHENKO K.A. 2002: *Melitaea cinxia aversa* ssp. n., a new subspecies of Nymphalidae from Tian-Shan (Lepidoptera, Rhopalocera). *Helios (Moscow)* **3**: 136–142.

- EHRlich P.R. & HANSKI I. 2004: *On the Wings of Checkerspots: A Model System for Population Biology*. Oxford University Press, New York, 371 pp.
- EXCOFFIER L., LAVAL G. & SCHNEIDER S. 2005: Arlequin ver. 3.0: An integrated software package for population genetics data analysis. *Evol. Bioinform. Online* **1**: 47–50.
- FOLMER O., BLACK M.B., HOCH W., LUTZ R.A. & VRIJHOCK R.C. 1994: DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol. Marine Biol. Biotechnol.* **3**: 294–299.
- FORISTER M.L., FORDYCE J.A. & SHAPIRO A.M. 2004: Geological barriers and restricted gene flow in the holarctic skipper *Hesperia comma* (Hesperiidae). *Mol. Ecol.* **13**: 3489–3499.
- GOLOBOFF P.A. 1998: *NONA. Program and Documentation*. Published by the author, New York.
- GÓMEZ A. & LUNT D.H. 2007: Refugia within refugia: patterns of phylogenetic concordance in the Iberian Peninsula. In Weiss S. & Ferrand N. (eds): *Phylogeography of Southern Europe Refugia*. Springer, London, in press.
- GRICHUK V.P. 1984: Late Pleistocene vegetation history. In Wright Jr.H.E. & Barnosky C.W. (eds): *Late Quaternary Environments of the Soviet Union*. University of Minnesota Press, Minneapolis, MN, pp. 155–178.
- HABEL J.C., SCHMITT T. & MÜLLER P. 2005: The fourth paradigm pattern of postglacial range expansion of European terrestrial species: the phylogeography of the Marbled White butterfly (Satyrinae, Lepidoptera). *J. Biogeogr.* **32**: 1489–1497.
- HANSKI I. 1999: *Metapopulation Ecology*. Oxford University Press, Oxford, 313 pp.
- HANSKI I. & KUUSSAARI M. 1995: Butterfly metapopulation dynamics. In Cappuccino N. & Price P.W. (eds): *Population Dynamics: New Approaches and Synthesis*. Academic Press, San Diego, CA, pp. 149–171.
- HANSKI I., KUUSSAARI M. & NIEMINEN M. 1994: Metapopulation structure and migration in the butterfly *Melitaea cinxia*. *Ecology* **75**: 747–762.
- HEWITT G.M. 1999: Post-glacial re-colonization of European biota. *Biol. J. Linn. Soc.* **68**: 87–112.
- HEWITT G.M. 2000: The genetic legacy of the Quaternary ice ages. *Nature* **405**: 907–913.
- HEWITT G.M. 2004: Genetic consequences of climatic oscillations in the Quaternary. *Philos. Trans. R. Soc. Lond. (B) (Biol. Sci.)* **359**: 183–195.
- HIGGINS L.G. 1941: An illustrated catalogue of the Palearctic *Melitaea* (Lep. Rhopalocera). *Trans. R. Entomol. Soc. Lond.* **91**: 175–365.
- IBRAHIM K.M., NICHOLS R.A. & HEWITT G.M. 1996: Spatial patterns of genetic variation generated by different forms of dispersal during range expansion. *Heredity* **77**: 282–291.
- JOYCE D. A. & PULLIN A.S. 2001: Phylogeography of the marsh fritillary *Euphydryas aurinia* (Lepidoptera: Nymphalidae) in the UK. *Biol. J. Linn. Soc.* **72**: 129–141.
- KUMAR S., TAMURA K. & NEI M. 2004: MEGA3: Integrated software for Molecular Evolutionary Genetics Analysis and sequence alignment. *Briefings Bioinform.* **5**: 150–163.
- KUUSSAARI M., SINGER M. & HANSKI I. 2000: Local specialization and landscape-level influence on host use in an herbivorous insect. *Ecology* **81**: 2177–2187.
- MICHAUX J.R., MAGNANOU E., PARADIS E., NIEBERDING C. & LIBOIS R. 2003: Mitochondrial phylogeography of the Woodmouse (*Apodemus sylvaticus*) in the Western Palearctic region. *Mol. Ecol.* **12**: 685–697.
- NIXON K.C. 2002: *Winclada*. Published by the author, Ithaca, NY.
- PAINTER J.N., SIITONEN J. & HANSKI I. 2007: Phylogeographical patterns and genetic diversity in three species of Eurasian boreal forest beetles. *Biol. J. Linn. Soc.* **91**: 267–279.
- SCHMITT T. 2007: Molecular biogeography of Europe: Pleistocene cycles and Postglacial trends. *Front. Zool.* **4**: 11.
- SCHMITT T. & SEITZ A. 2001: Intraspecific allozymatic differentiation reveals the glacial refugia and the postglacial expansions of European *Erebia medusa* (Lepidoptera: Nymphalidae). *Biol. J. Linn. Soc.* **74**: 429–458.
- SCHMITT T. & SEITZ A. 2002: Postglacial distribution area expansion of *Polyommatus coridon* (Lepidoptera: Lycaenidae) from its Ponto-Mediterranean glacial refugium. *Heredity* **89**: 20–26.
- SCHMITT T., RÖBER S. & SEITZ A. 2005: Is the last glaciation the only relevant event for the present genetic population structure of the meadow brown butterfly *Maniola jurtina* (Lepidoptera: Nymphalidae)? *Biol. J. Linn. Soc.* **85**: 419–431.
- TABERLET P., FUMAGALLI L., WUST-SAUCY A.-G. & COSSON J.-F. 1998: Comparative phylogeography and postglacial colonization routes in Europe. *Mol. Ecol.* **7**: 453–464.
- TARASOV P.E., VOLKOVA V.S., WEBB T., GUIOT J., ANDREEV A.A., BEZUSKO J.G., BYKOVA G.V., DUROFEYUK N.I., KVAVADZE E.V., OSIPOVA I.M., PANOVA N.K. & SEVASTYANOV D.V. 2000: Last glacial maximum biomes reconstructed from pollen and plant macrofossil data from northern Eurasia. *J. Biogeogr.* **27**: 609–620.
- TOLMAN T. & LEWINGTON R. 1997: *Butterflies of Britain and Europe*. HarperCollins Publishers, London, 320 pp.
- TUZOV V.K. 2000: *Guide to the Butterflies of Russia and Adjacent Territories. Vol. 2*. Pensoft, Sofia, 580 pp.
- WAHLBERG N. 2006: That awkward age for butterflies: insights from the age of the butterfly subfamily Nymphalinae. *Syst. Biol.* **55**: 703–714.
- WAHLBERG N. & ZIMMERMANN M. 2000: Pattern of phylogenetic relationships among members of the tribe Melitaeini (Lepidoptera: Nymphalidae) inferred from mtDNA sequences. *Cladistics* **16**: 347–363.
- WAHLBERG N., WEINGARTNER E. & NYLIN S. 2003: Towards a better understanding of the higher systematics of Nymphalidae (Lepidoptera: Papilionoidea). *Mol. Phylogenet. Evol.* **28**: 473–484.
- WAHLBERG N., BRABY M.F., BROWER A.V.Z., DE JONG R., LEE M.-M., NYLIN S., PIERCE N., SPERLING F.A., VILA R., WARREN A.D. & ZAKHAROV E. 2005a: Synergistic effects of combining morphological and molecular data in resolving the phylogeny of butterflies and skippers. *Proc. R. Soc. Lond. (B) (Biol. Sci.)* **272**: 1577–1586.
- WAHLBERG N., BROWER A.V.Z. & NYLIN S. 2005b: Phylogenetic relationships and historical biogeography of tribes and genera in the subfamily Nymphalinae (Lepidoptera: Nymphalidae). *Biol. J. Linn. Soc.* **86**: 227–251.
- VANDEWOESTIJNE S., BAGUETTE M., BRAKEFIELD P.M. & SACCHERI I.J. 2004: Phylogeography of *Aglais urticae* (Lepidoptera) based on DNA sequences of the mitochondrial COI gene and control region. *Mol. Phylogenet. Evol.* **31**: 630–646.
- WEINGARTNER E., WAHLBERG N. & NYLIN S. 2006: Speciation in Pararge (Satyrinae: Nymphalidae) butterflies – North Africa is the source of ancestral populations of all Pararge species. *Syst. Entomol.* **31**: 621–632.
- WIEMERS M. & FIEDLER K. 2007: Does the DNA barcoding gap exist? – a case study in blue butterflies (Lepidoptera: Lycaenidae). *Front. Zool.* **4**: 8.
- ZIMMERMANN M., WAHLBERG N. & DESCIMON H. 2000: Phylogeny of *Euphydryas checkerspot* butterflies (Lepidoptera: Nymphalidae) based on mitochondrial DNA sequence data. *Ann. Entomol. Soc. Am.* **93**: 347–355.

Received May 23, 2007; revised and accepted June 18, 2007